| Title | Page |
|---|------|
| I. Introduction | 1 |
| II. Literature Review | 4 |
| 1-Phytochemical studies. | 4 |
| 2. Toxicological studies. | 6 |
| 2.1 Insecticidal activit1y of some plant extracts | 6 |
| 2.1.1. Myzus persicae. | 6 |
| 2.1.2. Bemisia tabaci. | 19 |
| 2.2. Insecticidal activity of entomopathogenic fungi. | 36 |
| 2.2.1. Myzus persicae. | 36 |
| 2.2.2. Bemisia tabaci. | 39 |
| 3. Biological studies. | 44 |
| 4. Biochemical studies | 46 |
| 4.1. Biochemical effects of some plant extracts. | 46 |
| 4.2. Biochemical effects of fungicide. | 47 |
| 5. Susceptibility of some potato varieties crops against. | 47 |
| 5.1.Aphid Myzus persicae | 47 |
| 5.2. Bemisia tabaci. | 52 |
| 6. Effect of the chemical contents of potato leaves on | 53 |
| the population density of insect infestation. | |
| III-MATERIALS AND METHODS | 55 |
| 1. Rearing technique | 55 |
| 1.1. Rearing technique of potato aphid, <i>Myzus persicae</i> . | 55 |
| 1.2 Rearing technique of sweet potato whitefly, <i>Bemisia</i> | |
| tabaci. | |
| 2. Tested materials. | 57 |
| 2.1. Plant materials. | 57 |
| 2.1.1. Extraction technique of volatile oils (group A). | 59 |
| 2.1.1.1. Identification of volatile oils constituents | 59 |
| 2.1.2. Extraction technique of fixed oils (group B). | 60 |
| 2.1.2.1. Determination of free fatty acids. | 60 |
| 2.1.2.2. Isolation of unsaponifiable fractions | 60 |
| 2.2. Tested Biocide. | 60 |
| 3. Toxicological studies | 61 |
| 3.1. Preliminary insecticidal tests. | 61 |
| 3.2. Bioassay tests | 61 |
| 4. Biochemical studies. | 62 |

| 4.1. Preparation of sample for biochemical assays. | 62 | |
|--|-----|--|
| 4.2. Total proteins determination | | |
| 4.3. Total carbohydrates determination | | |
| 4.4. Total lipids determination | 64 | |
| 5. Susceptibility of certain potato varieties to the infestation | | |
| with aphids and whitefly. | | |
| 6. Effect of certain climatic factors (means of temperature and | | |
| relative humidity) on the population density of the dominant | | |
| homopterous insect infesting potato plant. | 65 | |
| 7. Effect of the chemical contents of potato leaves on the | 66 | |
| population density insects infestation | | |
| 7.1-Determination of protein content | 66 | |
| 7.2- Determination of carbohydrates content | 66 | |
| 7.3- Determination of phenol | 66 | |
| 7.4- Determination of PH value | 66 | |
| 7.5- Determination of K, P and N% | 66 | |
| Statistical analysis | 66 | |
| IV-RESULTS | 67 | |
| 1. Phytochemical Studies. | 67 | |
| 1.1. Extraction and identification of plant extracts. | 67 | |
| 1.1.1. Volatile oils. | | |
| 1.1.2. Fixed oils. | 102 | |
| 1.1.2.1. Identification of free fatty Acids. | 102 | |
| 1.1.2.2. Identification of the unsaponifiabla components. | | |
| 2. Toxicological studies. | 105 | |
| 2.1. Toxicological effects against adult of <i>M. persicae</i> and | 105 | |
| nymphs of <i>B. tabaci</i> under laboratory conditions. | | |
| 2.2. Susceptibility of Myzus persicae adult to the two volatile | | |
| oils. | | |
| 2.2.1. Susceptibility of adults of <i>Myzus persicae</i> and nymphs of | | |
| <i>B. tabaci</i> to the two fixed oils. | | |
| 2.2.2. Susceptibility of adults <i>Myzus persicae</i> and nymphs of | 112 | |
| <i>B. tabaci</i> to biozed. | | |
| 2.3. Effect of tested treatments at their LC_{50} on adult of <i>M</i> . | 114 | |
| persicae. | | |
| 3. Biological studies. | 115 | |
| - Biological activity of some plant treatments and one | 115 | |

| biocides against the adult of Myzus persicae. | |
|--|-----|
| 4. Biochemical studies. | |
| 4.1. The total protein. | |
| 4.2. The total lipids. | 121 |
| 4.3. The total carbohydrates. | 121 |
| 5. Ecological studies on certain piercing sucking insects and | 126 |
| their insect predators associated with different potato | |
| varieties cultivated in newly reclaimed sandy areas of El- | |
| Khattara district, Sharkia Governorate, Egypt . | |
| 5.1 Survey of insect pests and their predators associated with | 126 |
| potato varieties. | |
| 5.1.2. Population density of insect pests infesting potato | |
| vegetable plants | |
| 5.1.3. Population density of insect predators associated with | 127 |
| insect pests infesting potato vegetable plants. | |
| 5.2. Susceptibility of certain potato varieties to the infestation | |
| with aphids and whitefly. | |
| 5.2.1. Aphids (<i>Myzus persicae</i> and <i>Aphis gossypii</i>). | |
| 5.2.2. whitefly <i>B. tabaci</i> | |
| 5.3. Seasonal abundance of certain insect predators | |
| associated with piercing sucking insects infesting potato | |
| plants cultivated in newly reclaimed sandy areas at El- | |
| Khattara district, Sharkia Governorate, Egypt. | |
| 5.3.1. Chrysopids (<i>Chrysoperla carnea</i>). | 134 |
| 5.3.2. Coccinella septempunctata. | 135 |
| 5.3.3. Coccinella undecimpunctata. | 137 |
| 5.3.4. Metasyrphus corolla. | 139 |
| 5.3.5. Coccinella 9 pinctata. | 140 |
| 6. Effect of temperature and relative humidity on the | |
| population densities of aphids and whitefly infesting potato | |
| plants and their predators. | |
| 7. Effect of chemical contents of potato leaves on the | |
| population density of insects infestation. | |
| 1. Effect of protein content. | 143 |
| 1.1. Aphid insects. | 143 |
| 1.2. Whitefly insects. | 143 |

| 2. Effect of carbohydrate content. | |
|------------------------------------|-----|
| 2.1. Aphid insects. | 144 |
| 2.2. Whitefly insects. | 144 |
| 3. Effect of pH value. | 144 |
| 3.1. Aphid insects. | 145 |
| 3.2. Whitefly insects. | 145 |
| 4. Effect of total phenol. | |
| 4.1. Aphid insects. | 145 |
| 4.2. Whitefly insects. | 146 |
| V- Discussion | 148 |
| VI- Summary | 157 |
| VII- References | 163 |
| Arabic summary | 201 |

| Table | Title | Page |
|-------|--|------|
| No. | | |
| (1) | The plants used for screening of their insecticidal activities | |
| | against Myzus persicae and Bemisia tabaci. | 59 |
| (2) | GC/MS analysis of chemical constituents of Rosmarinus | |
| | officinalis volatile oil. | 69 |
| (3) | GC/MS analysis of chemical constituents of Ocimum | |
| | basilicum volatile oil | 83 |
| (4) | Free fatty acids and their relative percentage in Ricinus | |
| | communis and Nigella sativa oil analysis by gas-liquid | |
| | chromatography (GLC) | 103 |
| (5) | Identification the unsaponifiable matters of <i>Ricinus communis</i> | 105 |
| | and Nigella sativa oil analyzed by gas-liquid chromatography | |
| | (GLC). | |
| (6) | Toxicity of Ocimum basilicum and Rosmarinus officinalis | |
| | volatile oils against adult of <i>Myzus persicae</i> . | 106 |
| (7) | Toxicity of Ocimum basilicum and Rosmarinus officinalis | |
| | volatile oils against nymphs of <i>B. tabaci</i> . | 108 |
| (8) | Toxicity of <i>Ricinus communis</i> and <i>Nigella Sativa</i> fixed oils | |
| | against adult of <i>Myzus persicae</i> . | 110 |
| (9) | Toxicity of <i>Ricinus communis and Nigella Sativa</i> fixed oils | |
| | against nymphs of <i>B. tabaci</i> . | 111 |
| (10) | Toxicity of biozed aganist adult of Myzus persicae. | 112 |
| (11) | Toxicity of biozed against nymphs of B tabaci | 113 |
| () | Tomenty of ofolog against hympils of D. tabact. | |
| (12) | Effect of tested treatments on mortality of adult <i>M</i> . | |
| | persicae. | 114 |
| (13) | Effect of tested treatment at LC_{50} on some biological aspects | 117 |
| | against adult of Myzus persicae. | |
| (14) | Changes of total protein, total lipid and total carbohydrate | |
| | levels in the adult of <i>M. persicae</i> treated with the tested | |
| | treatments after 72 h. | 122 |
| (15) | Relative abundance of aphids & whitefly and the associated | |
| | insect predators collected from potato plants during 2016 | |
| | and 2017 seasons at El-Khattara district, Sharkia | |
| | Governorate, Egypt. | 128 |
| (16) | Total mean numbers of aphids and whitefly infested potato | |
| | cultivars during 2016 and 2017 seasons at El-Khattara | |
| | district, Sharkia Governorate, Egypt. | 130 |
| (17) | Weekly total number of certain insect pests infesting | |
| | different potato cultivars during winter plantation of 2016 at | |
| | El-Khattara district. Sharkia Governorate, Egypt.1 | 132 |

| (18) | Weekly total number of certain insect pests infesting | |
|------|--|-----|
| | different potato cultivars during winter plantation of 2017 at | |
| | El-Khattara district, Sharkia Governorate, Egypt. | 133 |
| (19) | Total number of certain insect predators per sample (30 | |
| | leaves / weekly) of different potato cultivars during winter | |
| | plantation of 2016 at El-Khattara district, Sharkia | |
| | Governorate, Egypt. | 136 |
| (20) | Total number of certain insect predators per sample (30 | |
| | leaves / weekly) of different potato cultivars during winter | |
| | plantation of 2017 at El-Khattara district, Sharkia | 100 |
| | Governorate, Egypt. | 138 |
| (21) | Simple correlation (r), partial regression (p) and explained | |
| | variance (E.V.) between mean temperature and mean | |
| | relative humidity and population density of aphid and | |
| | whitefly infesting potato plants and their predators in El- | |
| | khattara district during 2016 and 2017 seasons. | 145 |
| (22) | Effect of total proteins, chrbohydrate, PH values, phenol, | |
| | calcium, maghnsium, phosphorus and Potassium | |
| | percentage of potato cultivars on the population density of | |
| | aphids and whitefly. | 147 |

| Figure No. | Title | Page |
|---------------|---|------|
| (1) | GC/MS chromatogram of <i>Rosmarinus officinalis</i> . | 70 |
| (2) | GC/MS chromatogram of <i>Ocimum basilicum</i> . | 85 |
| (3) | GLC analysis of saponifiable components (fatty acids) of | 104 |
| | Ricinus communis fixed oil. | |
| (4) | GLC analysis of saponifiable components (fatty acids) of | 104 |
| | Nigella saliva fixed oil. | |
| (5) | Log. Conc.Mortality regression lines of the plant volatile oils | 107 |
| | against adult of Myzus persicae. | |
| (6) | Log. Conc. Mortality regression lines of the plant volatile oils | 108 |
| | against nymph of <i>B. tabaci</i> . | |
| (7) | Log. Conc. Mortality regression lines of the plant fixed oils | 110 |
| | against adult of Myzus persicae. | |
| | | 111 |
| (8) | Log. Conc. Mortality regression lines of the plant fixed oils | 111 |
| (0) | against nymphs of <i>B. tabaci</i> . | 110 |
| (9) | Log. Conc. Mortanty regression lines of biozed against | 112 |
| (10) | Myzus persicue. | 113 |
| (10) | nymphs of <i>B tabaci</i> | 115 |
| (11) | Effect of different treatments against adult <i>M</i> persical at | 115 |
| (11) | different periods (days). | 110 |
| | Effect of LC ₅₀ s concentrations of different treatments on | 118 |
| (12) | longevity of adult <i>Myzus persicae</i> . | |
| | | |
| (13) | Effect of LC ₅₀ s concentrations of different treatments on pre- | 119 |
| | parturition, parturition and post -parturition periods of | |
| | adult of <i>Myzus persicae</i> . | |
| (14) | Effect of LC_{50} concentrations of different treatments on | 120 |
| | number of nymphs per adult of <i>Myzus persicae</i> . | 100 |
| (15) | Total protein levels in <i>M. persicae</i> treated with the tested | 123 |
| | treatments after 72h. | |
| (16) | Changes % of total protein after 72h levels in <i>M. persicae</i> | 123 |
| | treated with the tested treatments from the untreated one after | _ |
| | 72h. | |
| (17) | Total lipid levels in <i>M. persicae</i> treated with the tested | 124 |
| | treatments after 72h. | |

LIST OF FIGURES

| (18) | Changes % of total lipid after 72h levels in <i>M. persicae</i> | 124 |
|------|--|-----|
| | 72h | |
| (19) | Total carbohydrate levels in <i>M. persicae</i> treated with the tested treatments after 72h. | 125 |
| (20) | Changes % of total carbohydrates after 72h levels in <i>M. persicae</i> treated with the tested treatments from the untreated one after 72h. | 125 |

List of plats

| 1 | Potato varieties and potato infestation by insects | 56 |
|---|--|----|
| 2 | Tested materials | 58 |