



Cairo University

Cairo University
Faculty of Veterinary Medicine
Department of Parasitology

Some studies about *Theileria* spp. among sheep in Egypt

A thesis Presented by
Asmaa Abd Elwadod Mohamed Hegab

B.V.Sc. Benha University (2004)

For master degree
(Parasitology)

Under supervision of

Dr. Magdy Mostafa Fahmy
Professor Emeritus of Parasitology
Faculty of Veterinary Medicine
Cairo University

Dr. Olfat Anter Mahdy
Professor of Parasitology
Faculty of Veterinary Medicine
Cairo University

Dr. Ahmed Anwar Wahba
Chief Researcher
Department of Parasitology
Animal Health Research Institute
Agricultural Research Center

2017



Cairo University

Cairo University
Faculty of Veterinary Medicine
Department of Parasitology

Supervision sheet

Dr. Magdy Mostafa Fahmy
Professor Emeritus of Parasitology
Faculty of Veterinary Medicine
Cairo University

Dr. Olfat Anter Mahdy
Professor of Parasitology
Faculty of Veterinary Medicine
Cairo University

Dr. Ahmed Anwar Wahba
Chief Researcher
Parasitology Department
Animal Health Research Institute
Agricultural Research Center

2017



Cairo University

Cairo University.
Faculty of Veterinary Medicine.
Department of Parasitology.

Name: Asmaa Abd Elwadod Hegab.

Nationality: Egyptian.

Date and place of birth: 16-10-1982 (Kaliobia Governorate).

Specialty: Parasitology.

Title of thesis: Some studies about *Theileria* spp. among sheep in Egypt.

Supervisors:

Dr. Magdy Mostafa Fahmy

Professor Emeritus of Parasitology, Faculty of Veterinary Medicine, Cairo University

Dr. Olfat Anter Mahdy

Professor of Parasitology, Faculty of Veterinary Medicine, Cairo University

Dr. Ahmed Anwar Wahba

Chief researcher, Parasitology Department , Animal Health Research Institute,
Agricultural Research Center

Abstract

The present study was carried out to investigate the accurate status of ovine *Theileria* infection in sheep in Giza governorate, Egypt during the period from May 2013 to April 2014. A total of 347 sheep blood samples (240 from different flocks and 107 from different slaughter houses) blood samples collected from both sex and different age groups. Giemsa stained blood smears examination by light microscope showed that 15.56% (54/347) were infected with *Theileria* spp. Statistical analysis using chi-square (χ^2) found no statistical significance among percent of *Theileria* infection in different age, sex, season and animal locations. Also, 86 tissue specimens were collected from slaughtered sheep each sample was divided in to two parts one for preparing impression smears stained with 10% Giemsa and the other part were fixed in 10% neutral buffered formalin, and used for histopathology. Schizont was more prevalent in lymph node impression smears (92.85%) than spleen and lung ones (81.81% and 36.36% respectively). Pathological examination revealed mainly depletion and necrosis of lymphocytes in inspected sheep lymph nodes and spleen. Also, sever destructions in the lung tissue (emphysema, collapse, congestion, pneumonia and infiltration with inflammatory cells) were observed. PCR applied for amplification of a fragment of the 18S ribosomal DNA on 5 positive and 10 negative samples by microscopic examination. 40% (6/15) of examined samples were infected. The PCR products subjected to RFLP assay for differentiation of various *Theileria* spp. By using HpaII restriction fermentase enzyme. The restriction enzyme differentiated *Theileria* spp. to *T. lestoquardi*, *T. ovis* and *T. annulata*. The present study concluded that the pathological alteration in examined organs indicated infection of the inspected sheep with the pathogenic type (*T. lestoquardi*). PCR RFLP is a diagnostic tool enabling direct, highly specific identification of *Theileria* spp. when compared with microscopic examination. Also, *Theileria* spp. that infects sheep in Egypt is polyphyletic origin.

Keywords: Sheep, *Theileria*, PCR RFLP, Egypt.

CONTENTS

Title	Page
INTRODUCTION	1
REVIEW OF LITERATURE	
I- Taxonomy of <i>Theileria</i> species.....	4
II- Importance of <i>Theileria</i> infection in sheep.....	4
III- Incidence rate of <i>Theileria</i> spp. infection in sheep.....	7
IV- Microscopic examination.....	16
V- Histopathological effect of <i>Theileria</i> spp. infection in sheep.....	20
VI- Polymerase Chain Reaction (PCR).....	22
VII- Restriction Fragment Length Polymorphism (RFLP).....	31
MATERIAL AND METHODS	
I- Investigated animals	34
II- Samples collection.....	34
III- Microscopic examination.....	35
IV- Statistical analysis.....	36
V- Histopathological examination.....	36
VI- Polymerase chain reaction (PCR).....	37
VII- Restriction fragment length polymorphism (RFLP) of PCR products.....	41
VIII- Agarose gel electrophoresis.....	43
RESULTS	
I-Incidence of <i>Theileria</i> spp. infection among sheep.....	45
Incidence of infection with <i>Theileria</i> spp. in inspected sheep from different sources.....	47
Incidence of infection with <i>Theileria</i> spp. among different exes.....	48
Incidence of infection with <i>Theileria</i> spp. in different age groups.....	49

II- Morphology of <i>Theileria</i> spp. in blood smears.....	50
Incidence of different stages of <i>Theileria</i> spp in blood smears.....	52
Microscopic examination of impression smears from slaughtered sheep internal organs.....	53
Incidence of <i>Theileria</i> spp. schizont in impression smears from sheep internal organs	54
Comparison between blood smear and impression smear for detection of <i>Theileria</i> infection.	55
III- Histopathological examination.....	56
IV- Polymerase chain reaction (PCR).....	61
V- RFLP of PCR products.....	63
DISCUSSION	65
CONCLUSION	75
SUMMARY	77
REFERENCES	80
ARABIC SUMMARY

LIST OF TABLES

Number	Table	Page
1	Sequence of primers used in the present study.	39
2	Reaction mixture used in PCR procedures.	40
3	Thermal profile used in PCR procedures.	40
4	Monthly and seasonal incidence rates of <i>Theileria</i> spp. infection among sheep in Giza governorate.	45
5	Incidence of <i>Theileria</i> infection in resident and preslaughtered sheep.	47
6	Incidence of <i>Theileria</i> spp. infection among different sheep sexes.	48
7	Incidence of <i>Theileria</i> infection among different age groups of sheep.	49
8	Incidence of different stages of <i>Theileria</i> spp. in positive sheep blood smears.	52
9	Incidence of <i>Theileria</i> spp. shizonts in impression smears from slaughtered sheep internal organs.	54
10	Comparison between different methods for the detection of <i>Theileria</i> infection.	55
11	Comparison between PCR and microscopic examination of <i>Theileria</i> spp. in sheep.	62
12	The pattern of RFLP of PCR products of different <i>Theileria</i> spp. by the use of HpaII.	64

LIST OF GRAPHS

Number	Graph	Page
1	Monthly incidence rates of <i>Theileria</i> spp. infection among sheep during the period from May 2013 to April 2014.	46
2	Seasonal incidence of <i>Theileria</i> Infection among sheep by microscopic examination.	46
3	Incidence of <i>Theileria</i> infection in resident and preslaughtered sheep.	47
4	Incidence of <i>Theileria</i> spp. infection among different sheep sexes.	48
5	Incidence of <i>Theileria</i> infection among different age groups of sheep.	49
6	Incidence of different stages of <i>Theileria</i> spp. in sheep blood.	52
7	Incidence of <i>Theileria</i> spp. shizonts Impression smears from slaughtered sheep internal organs.	55
8	Comparison between blood smear and impression smear for the detection of <i>Theileria</i> infection in sheep.	56
9	Comparison between PCR and microscopic examination of <i>Theileria</i> spp. in sheep.	62

LIST OF FIGURES

Number	FIGURE	Page
1	Giemsa-stained sheep blood smears, <i>Theileria</i> spp. erythrocytic form (X1000).	50
2	Giemsa-stained blood smears, <i>Theileria</i> spp. Koch's blue bodies (X1000).	51
3	Giemsa-stained impression smears from mesenteric lymph nodes showing <i>Theileria</i> spp. Koch's blue bodies (X1000).	53
4	Giemsa-stained impression smears from internal organ showing <i>Theileria</i> spp. Koch's blue bodies A: Microschizont in spleen B: Microschizont in lung (X1000).	54
5	Lymph node of sheep infected with <i>T. lestoquardi</i> showing infiltration to the fibrous capsule with leukocytes (HE) at X 400.	57

6	Lymph node of sheep infected with <i>T. lestoquardi</i> A: necrosis and depletion of lymphoid cells in a lymphoid follicle (X 200). B: Dilation of medullary sinuses with infiltration with histocytes, lymphocytes and lymphoblasts X 100 (HE). .	58
7	Spleen of sheep infected with <i>T. lestoquardi</i> . A. Depletion in the splenic follicle. B. Necrosis of lymphoid cells in the germinal center (HE) at X200.	59
8	Lung of sheep infected with <i>T. lestoquardi</i> . Showing emphysema (thin arrow) and collapse of adjacent alveoli (thick arrow) (HE) at X100.	60
9	Lung of sheep infected with <i>T. lestoquardi</i> . Showing thickening of alveolar wall (thin arrow) and privascular leukocytic aggregations (thick arrow) (HE) at X 200.	61
10	Agarose-gel electrophoresis of amplication products obtained from <i>Theileria</i> spp.	63
11	Restriction digests of <i>Theileria</i> sp. amplication products	64

LIST OF ABBREVIATIONS

Abbreviations	Description
IFA	Indirect Fluorescent Antibody
ME	Microscopic Examination
PCR	Polymerase Chain Reaction
RFLP	Restriction Fragment Length Polymorphism
EDTA	Ethylene Diamine Tetra Acetic Acid
MOT	malignant ovine theileriosis
PH	Degree of acidity and alkalinity
PBS	Phosphate buffer salin
μ l	Micro liter
%	Percent
DNA	Deoxy ribo Nuclic Acid
RPM	Revolution per minute
μ m	Micrometer
χ^2	Pearson chi-square
HE	Hematoxylin and eosin