



MANSOURA UNIVERSITY
FACULTY OF SCIENCE
BOTANY DEPARTMENT

Studies on the effect of using some nanomaterials in controlling stored grains pests

A Thesis

**Submitted in partial fulfillment of the requirements for the award of
the Master degree of Science in Botany
(Plant Physiology)**

By

Rehab Ibrahim Mohammed Gad

B.Sc. Botany, 2003

Botany Department

Faculty of Science (Girls)

Al-Azhar University

Cairo

Supervisors

Prof. Dr. Samia A. Haroun

Prof. of Plant Physiology

Botany Department

Faculty of Science - Mansoura University

Prof. Dr. Mahmoud E. El-Naggar

Prof. of Acarology

Plant Protection Research Institute

Agricultural Research Center

Dr. Doaa Mohamed Zein El-Abdein

Boraei

Assistant prof. Stored Products and Cereals

Research Department

Plant Protection Research Institute

Agricultural Research Center

2019

Contents

List of Tables.....	I
List of Figures.....	IV
List of abbreviations.....	VII
1.INTRODUCTION.....	1
Aim of the work.....	5
2. REVIEW OF LITERATURE.....	6
3. MATERIALS AND METHODS.....	30
Materials.....	30
Time course of experiments	30
Storage experiment.....	31
Practical methods	32
3.1. Transmission Electron Microscope (TEM).....	32
3. 2. Rearing	32
3.2.1. Mites rearing	32
3.2.2. Insects rearing.....	32
3.3. Bioassay tests	33
3.3.a. The insecticidal efficacy of zinc oxide and hydrophilic silica (NPs).....	33
Against target mites.....	33
Against target insects.....	34
Mortality percentage	35
3.4. Persistence tests (Storage).....	35
3.5 Tests on seeds stored for six months	36
3.5.1. Biochemical aspects of the seeds.....	36
3.5.2. Germination percentage and growth parameter.....	37
3.5.3. Estimation of photosynthetic pigments.....	38
3.5.4. Safety tests.....	39

3.5.4.1. Silicon estimation in wheat seedlings	39
3.5.4.2. Toxicity experiment of hydrophilic silica (NPs) on male mice.....	39
Data analysis.....	40
4. Experimental Results.....	41
4.1. Structure study.....	41
4.2. Insecticidal toxicity bioassay of <i>T. putrescentiae</i> and <i>C. berlesei</i>	41
4.2.1. Mortality percentage of <i>T. putrescentiae</i> adults on wheat grains	44
4.2.2. Insecticidal toxicity bioassay of <i>T. putrescentiae</i> by Abbot's method.....	47
4.2.3. Mortality percentage of <i>Caloglyphus berlesei</i> adults on wheat flour	50
4.2.4. Insecticidal toxicity bioassay of <i>C. berlesei</i> by Abbot's method.....	52
4.3. Insecticidal toxicity bioassay of <i>S. oryzae</i> , <i>C. maculatus</i> and <i>T. castaneum</i>	56
4.3.1.. Mortality percentage of rice weevil, <i>Sitophilus oryzae</i> adults on wheat	57
4.3.2. Insecticidal toxicity bioassay of <i>S. oryzae</i> by Abbot's method.....	58
4.3.3. Mortality percentage of cowpea beetle, <i>Callosobruchus maculatus</i> adults.....	63
4.3.4. Insecticidal toxicity bioassay of <i>C. maculatus</i> by Abbot's method.....	64
4.3.5. Mortality percentage of red flour beetle <i>Tribolium castaneum</i> adults	69
4.3.6. Insecticidal toxicity bioassay of <i>T. castaneum</i> by Abbot's method.....	70

4.4. Data analysis for mortality percentage of <i>Tyrophagus putrescentiae</i> , <i>Caloglyphus berlesei</i> on wheat flour, <i>Sitophilus oryzae</i> on wheat and <i>Callosobruchus maculatus</i> , on cowpea	75
4.5. Effect of LC90 of hydrophilic silica (NPs) against tested insects under storage conditions.....	78
4.5.1. Residual toxicity experiment.....	78
4.5.2. Wheat and cowpea components before and after storage for 6 months with hydrophilic silica (NPs).....	80
4.5.3. Changes in growth parameters of stored wheat grains treated with hydrophilic silica (NPs).....	82
4.5.4. Pigments content of hydrophilic silica-treated wheat seedlings for 6 months	84
4.5.5. Silicon content of wheat seedlings.....	85
4.6. Toxicity of hydrophilic silica (NPs) on male mice.....	86
4.6.1. Animal observations, food consumption, and changes in body weights.....	86
4.6.2. Histopathological Examinations.....	87
4.6.2.1. Liver histology.....	87
4.6.2.2. Lung histology.....	88
4.6.3. Biochemical examination.....	91
5. Discussion and conclusions.....	92
Insecticidal toxicity bioassay of <i>T. putrescentiae</i> and <i>C. berlesei</i>	93
Insecticidal toxicity bioassay of <i>S. oryzae</i> , <i>C. maculatus</i> and <i>T. castaneum</i>	94
Effect of LC90 of hydrophilic silica (NPs) against insects.....	98

Residual toxicity experiment.....	98
Component analysis of wheat grains and cowpea seeds.....	99
Changes in germination percentage and growth parameters of stored wheat grains.....	101
Chlorophyll and carotene content of wheat seedlings.....	102
Silicon residues in wheat seedlings	103
Toxicity of hydrophilic silica (NPs) on male mice.....	104
Liver and lung histology of mice.....	104
Biochemical evaluation.....	126
Summary.....	107
References.....	110
Arabic summary.....	1

List of Tables

Table (1) Mortality percentage of *Tyrophagus putrescentiae* adults on wheat flour treated with different levels of zinc oxide and hydrophilic silica nanoparticles during 7 days.....45

Table (2) Mortality percentage of *Tyrophagus putrescentiae* adults on wheat flour treated with different levels of zinc oxide and hydrophilic silica nanoparticles during 7 days by **Abbot's** method.....48

Table (3) Mortality percentage of *Caloglyphus berlesei* adults on wheat flour treated with different levels of zinc oxide and hydrophilic silica nanoparticles during 7 days.....51

Table (4) Mortality percentage of *Caloglyphus berlesei* adults on wheat flour treated with different levels of zinc oxide and hydrophilic silica nanoparticles during 7 days with **Abbot's** method.....54

Table (5) Mortality percentage of rice weevil, *Sitophilus oryzae* adults on wheat treated with different levels of zinc oxide and hydrophilic silica nanoparticles during 10 days.....59

Table (6) Mortality percentage (mean \pm SE) of *Sitophilus oryzae* adults (reared on stored wheat grains) treated with different levels of zinc oxide and hydrophilic silica nanoparticles throughout 10 days by **Abbot's** method.....61

Table (7) Mortality percentage of cowpea beetle, *Callosobruchus maculatus* adults treated with different levels of zinc oxide and hydrophilic silica nanoparticles during 5 days.....65

Table (8) Mortality percentage (mean \pm SE) of *Callosobruchus maculatus* adults (reared on stored cowpea seeds) treated with different levels of zinc oxide and hydrophilic silica nanoparticles throughout 5 days by **Abbot's** method.....67

Table (9) mortality percentage of red flour beetle *Tribolium castaneum* adults of wheat flour treated with different levels of zinc oxide and hydrophilic silica nanoparticles during 21 day.....71

Table (10) Mortality percentage (mean \pm SE) of *Tribolium castaneum* adults (reared on stored wheat flour) treated with different levels of zinc oxide and hydrophilic silica nanoparticles throughout 21 days by **Abbot's** method.....73

Table (11): Relative potency values of hydrophilic silica (NPs) on both *C. maculatus* and *S. oryzae* adults by **Probit** statics.....75

Table (12) Adult mortality percentage of *S. oryzae* and *C. maculatus* during 6 months storage of wheat grains and cowpea seeds treated with silica (NPs) (0.4g/kg).....79

Table (13) Wheat and cowpea components before and after storage for 6 months with hydrophilic silica (NPs).....81

Table (14) Parameters of germinated wheat after 6 months storage period of treated and untreated grains83

Table (15) Chlorophyll and carotene content of wheat seedlings after storage for 6 months84

Table (16) Silicon content in wheat seedlings before and after storage for 6 months after treatment with hydrophilic silica (NPs).....85

Table (17) Changes in body weight of control and treated mice exposed to hydrophilic silica (NPs) after 2 weeks of treatment and at the end of the experiment (4 weeks).....86

Table (18) Blood biochemical indices of the experimental mice feeding on wheat grains untreated and treated with hydrophilic silica (NPs).....91

List of Figures

Figure 1(A) Mites on (wheat flour), (B) *Sitophilus oryzae* on (wheat grains) (C) *Callosobruchus maculatus* on (cowpea seeds) (D) *Tribolium castaneum* on (wheat flour).....4

Figure 2. The TEM images of hydrophilic silica nanoparticles (x=40) X: magnification power.....41

Figure 3. The TEM images of zinc oxide nanoparticles (x=40) X: magnification power.....42

Figure 4. Mortality percentage of *Tyrophagus putrescentiae* adults on wheat flour treated with different levels of zinc oxide (A) and hydrophilic silica nanoparticles (B) during 7 days.....46

Figure 5. Mortality percentage of *Tyrophagus putrescentiae* adults on wheat flour treated with different levels of zinc oxide (A) and hydrophilic silica nanoparticles (B) during 7 days by **Abbot's** method.....49

Figure 6. Mortality percentage of *Caloglyphus berlesei* adults on wheat flour treated with different levels of zinc oxide (A) and hydrophilic silica nanoparticles (B) during 7 days.....53

Figure 7. Mortality percentage of *Caloglyphus berlesei* adults on wheat flour treated with different levels of zinc oxide (A) and hydrophilic silica nanoparticles (B) during 7 days with Abbot's method.....55

Figure 8. Mortality percentage of rice weevil, *Sitophilus oryzae* adults on wheat treated with different levels of zinc oxide (A) and hydrophilic silica nanoparticles (B) during 10 days.....60

Figure 9. Mortality percentage (mean \pm SE) of *Sitophilus oryzae* adults (reared on stored wheat grains) treated with different levels of zinc oxide (A) and hydrophilic silica nanoparticles (B) throughout 10 days by Abbot's method.....62

Figure 10. Mortality percentage of cowpea beetle, *Callosobruchus maculatus adults* treated with different levels of zinc oxide (A) and hydrophilic silica nanoparticles(B) during 5 days.....66

Figure 11. Mortality percentage (mean \pm SE) of *Callosobruchus maculatus* adults (reared on stored cowpea seeds) treated with different levels of zinc oxide (A) and hydrophilic silica nanoparticles (B) throughout 5 days by Abbot's method.....68

Figure 12.Mortality percentage of red flour beetle *Tribolium castaneum adults* on treated wheat flour with different levels of zinc oxide (A) and hydrophilic silica nanoparticles(B) during 21 days.....72

Figure 13. Mortality percentage (mean \pm SE) of *Tribolium castaneum* adults (reared on stored wheat flour) treated with different levels of zinc oxide (A) and hydrophilic silica nanoparticles (B) throughout 21 days by Abbot's method.....74

Figure 14(A). Toxicity lines of hydrophilic silica (NPs) against *C. maculatus*.....76

Figure 14(B). Toxicity lines of hydrophilic silica (NPs) against *S. oryzae*.....77

Figure (15) Germination of wheat stored for 6 months of treated and untreated grains.....83

Figure (16) Histological images of livers from 3 animals feeding on wheat grains untreated and treated with hydrophilic silica (NPs) after 28 days.....89

Figure (17)Histological images of Lungs from 3 animals feeding on wheat grains untreated and treated with hydrophilic silica (NPs) after 28 days.....90