

**ENVIRONMENTAL MANAGEMENT AND
BIOLOGICAL ASPECTS OF THE ERIOPHYID
MITES INFESTING SOME VEGETABLE CROPS**

By

BASSEM MAHMOUD MOHAMED FARAHAT

B. Sc. Agric., (Agric. Zoology and Nematology), Faculty of Agric., Cairo Al-
Azhar University, 2003.

M. Sc. Agric., (Plant Protection- “Acarology”), Faculty of Agric., Ain Shams
University, 2012.

THESIS

**Submitted in Partial Fulfillment of The
Requirements for the Degree**

Of

DOCTOR OF PHILOSOPHY

In

AGRICULTURAL SCIENCES

(Agric. Zoology and Nematology–“Acarology”)

**Department of Agricultural Zoology and Nematology
Faculty of Agriculture, Cairo
Al-Azhar University**

1441 A.H.

2020 A.D.

CONTENTS

Subject	Page
GENERAL INTRODUCTION	1-3
I. REVIEW OF LITERATURE	4
A- Effect of weather factors on mites populations.....	4-9
B- The role of predacious mites in reduction phytophagous mite populations.....	10-18
C- Incidence of phytophagous mites on horticultural crops...	19-29
D- Effect of some phytochemical components on the infestation rates of some injurious mites.	30- 37
E- Effect of anatomical characters and Trichomes types on the infestation of some injurious mites.	38-40
F- Effect of Different safe compounds for control some injurious mites.	41-55
G- Biological studies of phytophagous mites and predacious phytoseiid mites.	56-84
i. Eriophyids	56-65
ii. Phytoseiids	66-84
II. MATERIAL AND METHODS	85
A- Sampling and population assessment	85-86
B- Effect of Eriophyoid mites on Macro-Micronutrients and bio-chemicals contents of tomato, eggplant and artichoke cultivars leaves.	87-92
C- Anatomical characters (epidermal layer, cuticle thicknesses and trichomes types) of tomato, eggplant and artichoke cultivar.	93-94
D- Efficacy of recent acaricides for control eriophyid mites.	95-97
E- Biological studies	98-103
III- RESULTS AND DISCUSSIONS	104-
1- Seasonal fluctuations in phytophagous and predacious mite populations on different vegetable crops	104-139
A- Climatic conditions.....	106-107

Contents: continued

B- Population dynamics of the tomato russet mite and its phytoseiid predator on three tomato cultivars	108-139
i- Tomato 765 cultivar	112-122
ii- Safeera tomato cultivar	123-129
iii- Tomato 448 cultivar	130-139
C - Population dynamics of the egg-plant rust mite and its phytoseiid predator on three egg-plant cultivars:	140-170
i- Batra egg-plant cultivar	144-154
ii- N-Suma egg-plant cultivar	155-161
iii- Black beauty egg-plant cultivar	162-170
D - Population dynamics of the artichoke eriophyid mite on two artichoke cultivars	171
i- <u>Egyptian artichoke cultivar</u>	173
ii- <u>French artichoke cultivar</u>	184
2- Evaluation of the relationship between leaf photochemical components and harmful eriophyid mite infestations.	192
i- Tomato cultivars (765, Safeera and 448)...	194
ii- Eggplant cultivars (Batra, N-suma and Black beauty)	204
iii- Artichoke cultivars (Egyptian and French)	213

Contents: continued

3- Histological studies on the cuticle and lower epidermal layers of <i>Solanaceae</i> and <i>Asteraceae</i> cultivars.	225
i- Botanical observations	227
ii- Trichomes types	235
4- Efficacy of recent safe compounds for control of eriophyid tomato, eggplant and artichoke mites.	244
5- Biological studies	256
i- Eriophyid mites	256
1- The tomato russet mite, <i>A.lycopersici</i>	258
2- The eggplant rust mite, <i>A.melongenus</i>	282
3- The artichoke rust mite, <i>A.neocynarae</i>	304
ii- Phytoseiid mites:	325
1- <i>Amblyseius swirskii</i>	325
2- <i>Cydnoseius negevi</i>	337
SUMMARY	351-361
REFERENCES	362-420
ARABIC SUMMARY	1-11

Tables No.	Page
(1): Population dynamics of <i>A.lycopersici</i> and its predator <i>A.swirskii</i> on the tomato 765 cultivar during the first season (2016 / 2017)	118
(2): Population dynamics of <i>A.lycopersici</i> and its predator <i>A.swirskii</i> on the tomato 765 cultivar during the second season (2017 / 2018)	119
(3): Population dynamics of <i>A.lycopersici</i> and its predator <i>A.swirskii</i> on the Safeera tomato cultivar during the first season (2016 / 2017)	127
(4): Population dynamics of <i>A.lycopersici</i> and its predator <i>A.swirskii</i> on the Safeera tomato cultivar during the second season (2017 / 2018)	128
(5): Population dynamics of <i>A.lycopersici</i> and its predator <i>A.swirskii</i> on the tomato 448 cultivar during the first season (2016 / 2017)	134
(6): Population dynamics of <i>A.lycopersici</i> and its predator <i>A.swirskii</i> on the tomato 448 cultivar during the second season (2017 / 2018)	135
(7): Correlation coefficient between each of temperature, relative humidity and <i>A.lycopersici</i> population on the three tomato cultivars during two successive years (2016 / 2018).	137
(8): Population dynamics of <i>A.melongenus</i> and its predator <i>C.negevi</i> on the Batra egg-plant cultivar during the first season (2016 / 2017)	150
(9): Population dynamics of <i>A.melongenus</i> and its predator <i>C.negevi</i> on the Batra egg-plant cultivar during the second season (2017 / 2018)	151

Contents: continued

(10): Population dynamics of <i>A.melongenus</i> and its predator <i>C.negevi</i> on the N-Suma egg-plant cultivar during the first season (2016 / 2017)	159
(11): Population dynamics of <i>A.melongenus</i> and its predator <i>C.negevi</i> on the N-Suma egg-plant cultivar during the second season (2017 / 2018)	160
(12): Population dynamics of <i>A.melongenus</i> and its predator <i>C.negevi</i> on the Black beauty egg-plant cultivar during the first season (2016 / 2017)	167
(13): Population dynamics of <i>A.melongenus</i> and its predator <i>C.negevi</i> on the Black beauty egg-plant cultivar during the second season (2017 / 2018)	168
(14): Correlation coefficient between each of temperature, relative humidity and <i>A.melongenus</i> population on the three egg-plant cultivars during two successive years (2016 / 2018).	170
(15): Population dynamics of <i>A.neocynarae</i> on the Egyptian artichoke cultivar during the first season (2016 / 2017)	179
(16): Population dynamics of <i>A.neocynarae</i> on the Egyptian artichoke cultivar during the second season (2017 / 2018)	180
(17): Population dynamics of <i>A.neocynarae</i> on the French artichoke cultivar during the first season (2016 / 2017)	188
(18): Population dynamics of <i>A.neocynarae</i> on the French artichoke cultivar during the second season (2017 / 2018)	189
(19): Correlation coefficient between each of temperature, relative humidity and <i>A.neocynarae</i> population on the two artichoke cultivars during two successive years (2016 - 2018).	191

Contents: continued

(20): Average weight for twenty leaves of both healthy and heavily infested tomato cultivars (765, Safeera and 448) leaves by the tomato russet mite <i>Aculops lycopersici</i>	197
(21): Average macronutrient % and micronutrient ppm and bio-chemicals in healthy and heavily infested <i>Solanum lycopersici</i> L. leaves by the tomato russet mite <i>Aculops lycopersici</i> of different tomato cultivars.	198
(22): Correlation coefficient between macro - and micronutrients and <i>Aculops lycopersici</i> population on heavily infested leaves of different tomato cultivars.	199
(23): Average weight for twenty leaves of both healthy and heavily infested tomato cultivars (Batra, N-suma and Black beauty) leaves by the tomato russet mite <i>Aceria melongenensis</i>	206
(24): Average macronutrient % and micronutrient ppm and bio-chemicals in healthy and heavily infested <i>Solanum melongena</i> L. leaves by the eriophyid mite <i>Aceria melongenensis</i> of different egg-plant cultivars.	207
(25): Correlation coefficient between macro - and micronutrients and <i>Aceria melongenensis</i> population on heavily infested leaves of different egg-plant cultivars.	208
(26): Average weight for sixty discs of both healthy and heavily infested artichoke cultivars (Egyptian and French) leaves by the eriophyid mite <i>Aceria neocynarae</i>	218
(27): Average macronutrient % and micronutrient ppm and bio-chemicals in healthy and heavily infested <i>Cynara scolymus</i> L. leaves by the eriophyid mite <i>Aceria neocynarae</i> of different artichoke cultivars.	219
(28): Correlation coefficient between macro - and micronutrients and <i>Aceria neocynarae</i> population on heavily infested leaves of different two artichoke cultivars.	220
(29):Anatomical characters on cuticle and lower epidermal layers of different tomato cultivars infested with <i>A. lycopersici</i> mite	229

Contents: continued

(30): Anatomical characters on cuticle and lower epidermal layers of different eggplant cultivars infested with <i>A. melongenus</i> mite	230
(31): Anatomical characters on cuticle and lower epidermal layers of two artichoke cultivars infested with <i>A. neocynarae</i> mite	231
(32): Trichome types of the lower surfaces of the different tomato cultivars.	238
(33): Trichome types on the lower surfaces of the different eggplant cultivars	239
(34): Trichome types on the lower surfaces of the two artichoke cultivars.	239
(35): Acaricides effect on eriophyid tomato russet mite <i>Aculopslycopersici</i> at Behaira Gov. within 21 days after application treatments during 2017 season.	250
(36): Acaricides effect on eriophyid egg-plant mite <i>Aceria melongenus</i> at Behaira Gov. within 21 days after application treatments during 2017 season.	251
(37): Acaricides effect on eriophyid artichoke mite <i>Aceria neocynarae</i> at Behaira Gov. within 21 days after application treatments during 2018 season.	252
(38). Duration (in days) of Different stages and oviposition rate of <i>Aculops lycopersici</i> surviving on leaves of three tomato cultivars (765, Safeera and 448) at constant temperatures and 45 % R. H	260
(39). Relationship between temperature and developmental rate of the tomato russet mite, <i>Aculops lycopersici</i> (Massee) females on three tomato cultivars (45% RH).	263
(40). Relationship between temperature and developmental rate of the tomato russet mite, <i>Aculops lycopersici</i> (Massee) males on three tomato cultivars (45% RH).	268

Contents: continued

(41). Life table parameters of the tomato russet mite <i>Aculops lycopersici</i> survived on leaves of three tomato cultivars (765, Safeera and 448) at constant temperatures and 45 % R. H.	273
(42). Qualitative analysis of the spermatophores of the tomato russet mite <i>A.lycopersici</i> on three different tomato cultivars.	281
(43). Duration (in days) of Different stages and oviposition rate of <i>Aceria melongenus</i> surviving on leaves of three eggplant cultivars (Batra, N-suma and Black beauty) at three constant temperatures and 45 % R. H.	285
(44). Qualitative analysis of the spermatophores of the eggplant rust mite <i>A.melongenus</i> on three different eggplant cultivars.	287
(45). Relationship between temperature and developmental rate eggplant rust mite <i>Aceria melongenus</i> females on three eggplant cultivars (45% RH).	290
(46). Relationship between temperature and developmental rate of eggplant rust mite <i>Aceria melongenus</i> males on three eggplant cultivars (45% RH).	294
(47). Life table parameters of the eggplant rust mite <i>Aceria melongenus</i> survived on leaves of three eggplant cultivars (Batra, N-suma and Black beauty) at three constant temperatures and 45 % R. H.	300
(48). Duration (in days) of Different stages and oviposition rate of <i>Aceria neocynarae</i> surviving on leaves of two artichoke cultivars (Egyptian and French) at three constant temperatures and 45 % R. H.	307
(49). Qualitative analysis of the spermatophores of the artichoke rust mite <i>A.neocynarae</i> on two artichoke cultivars	310

Contents: continued

(50). Relationship between temperature and developmental rate of artichoke rust mite <i>Aceria neocynarae</i> females on two artichoke cultivars (45% RH).	314
(51). Relationship between temperature and developmental rate of artichoke rust mite <i>Aceria neocynarae</i> males on two artichoke cultivars (45% RH).	317
(52). Life table parameters of the artichoke rust mite <i>Aceria neocynarae</i> survived on leaves of two artichoke cultivars (Egyptian and French) at three constant temperatures and 45 % R. H.	321
(53).Developmental times (days) of the predatory phytoseiid mite, <i>Amblyseius swirskii</i> reared on tomato russet mite, <i>A.lycopersici</i> at different constant temperatures and 45% R.H.	327
(54).Relationship between temperature and developmental rate of <i>A. swirskii</i> females reared on tomato russet mite <i>A.lycopersici</i> (45% R. H.).	329
(55). Relationship between temperature and developmental rate of <i>A. swirskii</i> males reared on tomato russet mite <i>A.lycopersici</i> (45% R. H.).	331
(56). Life table parameters of the <i>A.swirskii</i> fed on the tomato russet mite, <i>A.lycopersici</i> at constant temperatures and 45%R.H.	335
(57).Developmental times (days) of the predatory phytoseiid mite, <i>Cydnoseius negevi</i> reared on eggplant rust mite, <i>A.melongenus</i> at different constant temperatures and 45% R.H.	341
(58). Life table parameters of the <i>C.negevi</i> fed on the eggplant rust mite, <i>A.melongenus</i> at constant temperatures and 45%R.H.	344

Contents: continued

(59). Relationship between temperature and developmental rate of <i>C. negevi</i> females reared on eggplant rust mite, <i>A.melongenus</i> (45% R. H.).	347
(60). Relationship between temperature and developmental rate of <i>C. negevi</i> males reared on eggplant rust mite, <i>A.melongenus</i> (45% R. H.).	349

Figures No.	Page
(1): Tomato russet mite, <i>Aculops lycopersici</i> (Masse) (After Abou-Awad, 1976)	110
(2): Symptoms of eriophyid mite infestation on tomato 765 cultivar	114
(3): Population trends of the tomato russet mite <i>A.lycopersici</i> and its predatory mite, <i>A.swirskii</i> associated with tomato cultivars in relation to temperature and relative humidity during two successive seasons (2016 / 2018)	120
(4): Percentage of Immature stages in the total population of the tomato russet mite <i>A.lycopersici</i> on tomato 765 cultivar during two successful seasons (2016 – 2018)	122
(5): Symptoms of eriophyid mite infestation on Safeera tomato cultivar	125
(6): Percentage of Immature stages in the total population of the tomato russet mite <i>A.lycopersici</i> on Safeera tomato cultivar during two successful seasons (2016 – 2018)	129
(7): Symptoms of eriophyid mite infestation on tomato 765 cultivar	132
(8): Percentage of Immature stages in the total population of the tomato russet mite <i>A.lycopersici</i> on tomato 448 cultivar during two successful seasons (2016 – 2018)	136
(9): Egg-plant rust mite, <i>Aceria melongenae</i> (Zaher & Abou-Awad)(After Zaher & Abou-Awad, 1978 / 1979)	142
(10): Symptoms of eriophyid mite infestation on Batra Egg-plant cultivar	146

Contents: continued

(11): Population trends of the egg-plant mite <i>A.melongenus</i> and its predatory mite, <i>C.negavi</i> associated with egg-plant cultivars in relation to temperature and relative humidity during two successive seasons (2016 / 2018)	152
(12): Percentage of Immature stages in the total population of the egg-plant rust mite <i>A.melongenus</i> on Batra cultivar during two successful seasons (2016 – 2018)	154
(13): Symptoms of eriophyid mite infestation on N-Suma Egg-plant cultivar	157
(14): Percentage of Immature stages in the total population of the egg-plant rust mite <i>A.melongenus</i> on N-Suma cultivar during two successful seasons (2016 – 2018)	161
(15): Symptoms of eriophyid mite infestation on Black beauty Egg-plant cultivar	165
(16): Percentage of Immature stages in the total population of the egg-plant rust mite <i>A.melongenus</i> on Black beauty cultivar during two successful seasons (2016 – 2018)	169
(17): Artichoke eriophyid mite, <i>Aceria neocynarae</i> (Keifer) (After Abou-Awad, 1976)	175
(18): Symptoms of eriophyid mite infestation on Egyptian Artichoke cultivar	177
(19): Population trends of the artichoke eriophyid mite <i>A.neocynarae</i> associated with artichoke cultivars in relation to temperature and relative humidity during two successive seasons (2016 / 2018)	181

Contents: continued

(20): Percentage of Immature stages in the total population of the artichoke eriophyid mite <i>A.neocynarae</i> on Egyptian artichoke cultivar during two successful seasons (2016 – 2018)	183
(21): Symptoms of eriophyid mite infestation on French Artichoke cultivar	186
(22): Percentage of Immature stages in the total population of the artichoke eriophyid mite <i>A.neocynarae</i> on French artichoke cultivar during two successful seasons (2016 – 2018)	190
(23):Growth parameters for both healthy and heavily infested tomato cultivar leaves (765, Safeera and 448) by the tomato russet mite, <i>A.lycopersici</i>	200
(24): Macronutrients (N, P, K, Ca and Mg) % and total proteins in both healthy and heavily infested tomato cultivar leaves (765, Safeera and 448) by the tomato russet mite, <i>A.lycopersici</i>	201
(25): Micronutrients (Fe, Zn, Mn and Cu) ppm in both healthy and heavily infested tomato cultivars leaves (765, Safeera and 448) by the tomato russet mite, <i>A.lycopersici</i>	202
(26):Total phenols, sugars and carbohydrates in both healthy and heavily infested tomato cultivar leaves (765, Safeera and 448) by the tomato russet mite, <i>A.lycopersici</i>	203
(27): Growth parameters for both healthy and heavily infested egg-plant cultivar leaves (Batra, N-Suma and 165Black beauty) by the egg-plant rust mite, <i>A.melongenus</i>	209

Contents: continued

(28): Macronutrients (N, P, K, Ca, Mg) % and total proteins in both healthy and heavily infested egg-plant cultivar leaves (Batra, N-Suma and Black beauty) by the egg-plant rust mite, <i>A.melongenus</i>	210
(29): Micronutrients (Fe, Zn, Mn and Cu) ppm in both healthy and heavily infested egg-plant cultivar leaves (Batra, N-Suma and Black beauty) by the egg-plant rust mite, <i>A.melongenus</i>	211
(30): Total phenols, sugars and carbohydrates in both healthy and heavily infested egg-plant cultivar leaves (Batra, N-Suma and Black beauty) by the egg-plant rust mite, <i>A.melongenus</i>	212
(31):Growth parameters for both healthy and heavily infested artichoke cultivars leaves (Egyptian and French) by the artichoke rust mite, <i>A.neocynarae</i>	221
(32): Macronutrients (N, P, K, Ca , Mg) % and total proteins in both healthy and heavily infested artichoke cultivars leaves (Egyptian and French) by the artichoke rust mite, <i>A. neocynarae</i>	222
(33): Micronutrients (Fe, Zn, Mn and Cu) ppm in both healthy and heavily infested artichoke cultivars leaves (Egyptian and French) by the artichoke rust mite, <i>A. neocynarae</i>	223
(34): Total phenols, sugars and carbohydrates in both healthy and heavily infested artichoke cultivars leaves (Egyptian and French) by the artichoke rust mite, <i>A. neocynarae</i>	224

Contents: continued

(35): C =Cuticle layer, E = Epidermal layer and L.= Leaf Blade of tomato cultivars.	232
(36): C =Cuticle layer, E = Epidermal layer and L.= Leaf Blade of eggplant cultivars.	233
(37): C =Cuticle layer, E = Epidermal layer and L.= Leaf Blade of artichoke cultivars.	234
(38): Trichome types of the lower surfaces (A-C)	240
(39): Trichome types of the lower surfaces (A-C)	241
(40): Trichome types of the lower surfaces (A-C)	242
(41): Trichome types of the lower surfaces (A-C)	243
(42): Number of tomato russet mite <i>A.lycopersici</i> in treated tomato 765 leaves at Behaira Governorate in 2017 season	253
(43): Number of eriophyid egg-plant russet mite <i>A.melongenus</i> in treated Batra leaves at Behaira Governorate in 2017 season.	254
(44): Number of eriophyid artichoke russet mite <i>A.neocynarae</i> in treated Egyptian leaves at Behaira Governorate in 2018 season.	255
(45). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. lycopersici</i> (Massee) females reared on tomato-765 cultivar	265
(46). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. lycopersici</i> (Massee) females reared on Safeera – tomato cultivar.	266
(47). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. lycopersici</i> (Massee) females reared on tomato-448 cultivar	267

Contents: continued

(48). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. lycopersici</i> (Massee) males reared on tomato-765 cultivar.	269
(49). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. lycopersici</i> (Massee) males reared on Safeera – tomato cultivar.	270
(50). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. lycopersici</i> (Massee) males reared on tomato-448 cultivar.	271
(51): Age-specific fecundity (Mx) and survivorship (Lx) of <i>A.lycopersici</i> fed on tomato 765, cultivar at three constant temperatures degree.	274
(52): Age-specific fecundity (Mx) and survivorship (Lx) of <i>A.lycopersici</i> fed on Safeera tomato cultivar at three constant temperatures degree.	275
(53): Age-specific fecundity (Mx) and survivorship (Lx) of <i>A.lycopersici</i> fed on tomato 448, cultivar at three constant temperatures degree.	276
(54). Spermatophores of the tomato russet mite <i>A.lycopersici</i> on tomato leaves.	280
(55). Spermatophores of the eggplant rust mite <i>A.melongenus</i> on tomato leaves.	286
(56). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. melongenus</i> females reared on eggplant Batra cultivar	291
(57). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. melongenus</i> females reared on eggplant N-suma cultivar	292
(58). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. melongenus</i> females reared on eggplant Black beauty cultivar	293
(59). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. melongenus</i> males reared on eggplant Batra cultivar	295

Contents: continued

(60). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. melongenus</i> males reared on eggplant N-suma cultivar.	296
(61). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. melongenus</i> males reared on eggplant Black beauty cultivar.	297
(62): Age-specific fecundity (Mx) and survivorship (Lx) of <i>A.melongenus</i> fed on Batra eggplant cultivar at three constant temperatures degree.	301
(63): Age-specific fecundity (Mx) and survivorship (Lx) of <i>A.melongenus</i> fed on N-suma eggplant cultivar at three constant temperatures degree.	302
(64): Age-specific fecundity (Mx) and survivorship (Lx) of <i>A.melongenus</i> fed on Black beauty eggplant cultivar at three constant temperatures degree.	303
(65). Spermatophores of the artichoke rust mite <i>A.neocynarae</i> on artichoke leaves.	311
(66). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. neocynarae</i> females reared on artichoke Egyptian cultivar	315
(67). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. neocynarae</i> females reared on artichoke French cultivar	316
(68). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. neocynarae</i> males reared on artichoke Egyptian cultivar	318
(69). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. neocynarae</i> males reared on artichoke French cultivar	319
(70): Age-specific fecundity (Mx) and survivorship (Lx) of <i>A.neocynarae</i> fed on Egyptian artichoke cultivar at three constant temperatures degree.	322

Contents: continued

(71): Age-specific fecundity (Mx) and survivorship (Lx) of <i>A.neocynarae</i> fed on French artichoke cultivar at three constant temperatures degree.	323
(72). <i>Amblyseius swirskii</i> , Athias – Henriot	326
(73). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. swirskii</i> females fed on tomato russet mite <i>A.lycopersici</i>	330
(74). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>A. swirskii</i> males fed on tomato russet mite <i>A.lycopersici</i>	332
(75): Age-specific fecundity (Mx) and survivorship (Lx) of <i>A.swirskii</i> on <i>A. lycopersici</i> at three constant temperatures degree.	336
(76): <i>Cydnoseius negevi</i> (Swirski & Amitai)	339
(77): Age-specific fecundity (Mx) and survivorship (Lx) of <i>A.swirskii</i> on <i>C. negevi</i> at three constant temperatures degree.	345
(78). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>C. negevi</i> females fed on eggplant rust mite <i>A.melongenus</i> .	348
(79). Linear regression analysis of temperature versus developmental rate, degree – day's requirements, and minimum developmental thresholds of <i>C. negevi</i> males fed on eggplant rust mite <i>A.melongenus</i> .	350