



*Faculty of Science
Entomology Department*

**Ecological and biological studies on aphids and their
associated predatory spiders on two field crops at
Qalubia governorate, Egypt.**

A Thesis

*Submitted for the degree of Ph. D. in Entomology
Presented to the Faculty of Science
Ain Shams University*

BY

Manal Husseiny Mohammed Ibraheem

B.Sc. Entomology (2005)

M.Sc. Entomology (2016)

Supervised by

Prof. Dr. Nawal Mahmoud Shanbaky

Prof. Dr. Nadia Helmy Ahmed

Prof. Dr. Mohammed Hassan El-Erksousy

Prof. Dr. Amal Ibraheem Abo-Zaed

Dr. Ayat Yousery Mohamed

2021

Abstract

The seasonal dynamics of the cowpea aphid, *Aphis craccivora* (Koch) and the corn leaf aphid, *Rhopalosiphum maidis* (Fitch) on the broad bean and maize crop, respectively, were investigated at Qaha research Station, Qalubya governorate, Egypt during two successive years, 2017 and 2018. Infestation of broad bean and maize started in January and May or June, respectively. The monthly collected number and percent abundance of most movable life stages gradually increased to reach peaks in March and August then decreased in the next month at the end of the season of each of the broad bean and maize, respectively. Immatures were the most dominant life stages collected in each month than adults, followed by unwinged and winged adult females descendingly but no adult males were collected throughout the broad bean and maize seasons in each year. Generally, the total collected number of each life stage of *A. craccivora* and *R. maidis* infested each of the broad bean and maize was higher in 2018 than 2017 and on broad bean than maize in each year.

Twenty six spider species belonging to 20 genera and 10 families of order Araneae were associated with the two aphids, *A. craccivora* and *R. maidis* and were collected on broad bean and maize crops, at Qaha research station in Qalubya governorate, Egypt during the two successive years, 2017 and 2018. During the

period of study, Philodromidae was the most abundant family of the total collected spiders (1486) on the two crops followed by Salticidae, Theridiidae, Thomisidae then Lycosidae followed by Cheiracanthidae and Araneidae, Linyphiidae, Dictynidae and Aglenidae.

On maize, the collected families did not include Dictynidae and Aglenidae in both years of study. The numbers of the monthly and total collected spiders of each family and species, percent occurrence and hence abundance of each family and species varied on the studied crops in each season. However, nearly 50% of the collected spider genera and species were common associates of both *A. craccivora* and *R. maidis* on broad beans and maize, respectively in one or both years of study.

Generally, in parallel with aphids on each of broad bean and maize, most of the spider families and the total number of collected spiders and some species started appearance with relatively small numbers in winter (January) and late spring to early summer (May and June) then gradually increased to reach peaks in March and August, then decreased at the end of each season in April and September, respectively. The total number of spiders collected was higher in 2018 than 2017 and on broad bean (421&357) than maize (362&346) in each year of study.

The effect of temperature and prey quality of aphid species life stage and their basic nutritional components on food consumption, predation rate and duration of development of spiderlings of *Thanatus albini* reared on a monotypic diet of immatures or adult females of each of *Aphis craccivora* and *Rhopalosiphum maidis* were studied at 3 different temperatures of 15, 27 and $35\pm 2^{\circ}\text{C}$ and RH of 60-70%. Also, effects of the prey quality of the monotypic diet of each aphid on the tested parameters were studied in spiderlings of *Thomisus spinifer* and on reproduction in adult females of both spider species at 27°C .

Of the three tested temperatures, 27°C was considered as the optimum temperature for aphid consumption and predation rate of spiderlings of *Th. albini* when fed on each of the two aphid species, due to increasing the two parameters and prolonging spiderling durations. Also, life, development of spiderlings and female reproduction of each of *Th. albini* and *T. spinifer* were maintained and completed at the optimum temperature of 27 by feeding on immatures or adult females' aphid (except the first three spiderlings) of each of *A. craccivora* and *R. maidis* in *Th. albini* and only of *R. maidis* in *T. spinifer*.

The two spider species showed efficacy as predators, consuming higher numbers of immatures than adults of each of the two aphid species with more consumption of *A. craccivora* than *R. maidis* in *Th. albini* and only of *R. maidis* in *T. spinifer* with the refusal of feeding on *A. craccivora*. The number of

consumed aphids by each spider species was increased by increasing the age of the spiderling stage.

Quantitative analysis of the basic nutritional components showed a higher concentration of total protein and carbohydrate in adult females than immatures of each aphid species but no significant difference between each of the immatures and adult females of the two aphid species. The significance of the obtained results was discussed.

Keywords:

Aphis craccivora, *Rhopalosiphum maidis*, Seasonal dynamics, Spiders, Survey

CONTENTS

Content	Page
I. INTROUDUCTION	1
II. LITERATURE REVIEW	4
1. Seasonal dynamics of <i>Aphis craccivora</i> and <i>Rhopalosiphum maidis</i>	4
2. Seasonal abundance of predatory spiders	16
3. Predation of spiders	27
4. Biochemical studies on <i>Aphis craccivora</i> and <i>Rhopalosiphum maidis</i>	38
III. MATERIALS AND METHODS	42
1. Ecological studies	42
1.1 Collection method of aphids	42
1.2 Collection method of associated predatory spiders	43
2. Biological studies	44
2.1 Aphid colonies	44
2.2 Spider laboratory colony	44
2.3 Determination of the predation rate of immatures and adult female spiders on adult and immatures	45

Content	Page
<i>Aphis craccivora</i> and <i>Rhopalosiphum maidis</i>	
3. Biochemical studies	46
3.1 Determination of total carbohydrates	46
3.2 Determination of total proteins	47
3.3 Determination of total lipids	48
4. Statistical analysis	48
IV. RESULTS	49
A. Ecological studies on Aphids	49
1. Seasonal dynamics of the cowpea aphid, <i>Aphis craccivora</i> infesting broad bean crop at Qalubya governorate	49
1.1 Seasonal dynamics of the cowpea aphid, <i>Aphis craccivora</i> during the broad bean season 2017	49
1.2 Seasonal dynamics of the cowpea aphid, <i>Aphis craccivora</i> during the broad bean season 2018	56
2. Seasonal dynamics of the corn leaf aphid, <i>Rhopalosiphum maidis</i> infesting maize crop at Qalubya governorate	63
2.1 Seasonal dynamics of the corn leaf aphid, <i>Rhopalosiphum maidis</i> infesting maize crop during 2017 season	63

Content	Page
2.2 Seasonal dynamics of the corn leaf aphid, <i>Rhopalosiphum maidis</i> infesting maize crop during 2018 season	70
B. Spider families associated with <i>Aphis craccivora</i> and <i>Rhopalosiphum maidis</i> on broad bean and maize crops	77
1. Spider families associated with <i>Aphis craccivora</i> on broad bean	77
2. Spider families associated with <i>Rhopalosiphum maidis</i> on maize crops	86
3. Population density of spiders and their prey of <i>Aphis craccivora</i> and <i>Rhopalosiphum maidis</i> on broad bean and maize	94
4. Survey and abundance of spider species of the collected families associated with <i>Aphis craccivora</i> and <i>Rhopalosiphum maidis</i> on broad beans and maize	102
C. Food consumption and predation rate of life stages of spiders fed on <i>Aphis craccivora</i> and <i>Rhopalosiphum maidis</i>	108
1. Effect of temperature and prey quality on	108

Content	Page
duration, food consumption and predation rate of spiderlings of <i>Thanatus albini</i> fed on immatures and adult females of <i>A. craccivora</i> and <i>R. maidis</i>	
2. Effect of Prey Quality on Duration, Food Consumption and Predation Rate of Spiderlings of <i>Thomisus spinifer</i> Fed on Immatures and Adult Females of <i>Rhopalosiphum maidis</i> at 27°C	129
3. Effect of prey quality on reproduction of female <i>Thanatus albini</i> reared on immatures of <i>Aphis craccivora</i> and <i>Rhopalosiphum maidis</i>	135
4. Effect of prey quality on reproduction of female <i>Thomisus spinifer</i> reared on immatures of <i>Rhopalosiphum maidis</i> .	137
D. Quantitative analysis of basic nutritional components of immatures and adult females of <i>Aphis craccivora</i> and <i>Ropalosiphum maidis</i>	139
V. DISCUSSION	144
VI. SUMMARY	159
VII. REFERENCES	166
ARABIC SUMMARY	
ARABIC ABSTRACT	

LIST OF TABLES

No.	Table title	Page
1	Seasonal dynamics of <i>A. craccivora</i> infesting broad bean crop during 2017 at Qaha station, Qalubya governorate.	52
2	Mean temperature and relative humidity at Qaha station, Qalubya governorate during the broad bean season 2017.	53
3	Seasonal dynamics of <i>A. craccivora</i> infesting broad bean crop during 2018 at Qaha station, Qalubya governorate.	59
4	Mean temperature and relative humidity at Qaha station, Qalubya governorate during the broad bean season 2018.	60
5	Seasonal dynamics of <i>Rhopalosiphum maidis</i> infesting maize crop during 2017 at Qaha station, Qalubya governorate.	66
6	Mean temperature and relative humidity at Qaha station, Qalubya governorate during the maize season 2017.	67
7	Seasonal dynamics of <i>Rhopalosiphum maidis</i> infesting maize crop during 2018 at Qaha station, Qalubya governorate.	73
8	Mean temperature and relative humidity at Qaha station, Qalubya governorate during the maize season 2018.	74

No.	Table title	Page
9	Number of collected spiders and percent occurrence of spider families on broad bean crop at Qaha station, Qalubya governorate during two successive years 2017 and 2018	80
10	Seasonal dynamics of spider families on broad bean crop at Qaha station, Qalubya governorate during 2017 season.	82
11	Seasonal dynamics of spider families on broad bean crop at Qaha station, Qalubya governorate during 2018 season.	84
12	Number of collected spiders and percent occurrence of spider families on maize crop at Qaha station, Qalubya governorate during two successive years 2017 and 2018.	88
13	Seasonal dynamics of spider families on maize crop at Qaha station, Qalubya governorate during 2017 season.	90
14	Seasonal dynamics of spider families on maize crop at Qaha station, Qalubya governorate during 2018 season	92
15	Population density of spiders and their prey of <i>Aphis craccivora</i> on broad bean crop (40 plants/month) at Qaha station, Qalubya governorate during two successive years 2017 and 2018.	96
16	Population density of spiders and their prey of <i>Rhopalosiphum maidis</i> on maize crop (40plants/month) at Qaha station, Qalubya governorate during two successive years 2017 and 2018.	99

No.	Table title	Page
17	Survey and collected number of spider species on broad bean crop at Qaha station, Qalubya governorate during 2017 season.	104
18	Survey and collected number of spider species on broad bean crop at Qaha station, Qalubya governorate during 2018 season.	105
19	Survey and collected number of spider species on maize crop at Qaha station, Qalubya governorate during 2017 season.	106
20	Survey and collected number of spider species on maize crop at Qaha station, Qalubya governorate during 2018 season.	107
21	Duration, food consumption and predation rate of spiderling stages of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Aphis craccivora</i> at 27± 2°C.	113
22	Duration, food consumption and predation rate of spiderling stages of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Aphis craccivora</i> at 35± 2°C.	117
23	Duration, food consumption and predation rate of spiderling stages of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 27± 2°C.	121

No.	Table title	Page
24	Duration, food consumption and predation rate of spiderling stages of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 35± 2°C.	125
25	Duration, food consumption and predation rate of spiderling stages of <i>Thomisus spinifer</i> (F: Thomisidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 27± 2°C.	131
26	Biological aspects of reproduction of adult female <i>Thanatus albini</i> reared on immatures of <i>Aphis craccivora</i> and <i>Rhopalosiphum maidis</i> at 27± 2°C.	136
27	Biological aspects of reproduction of adult female <i>Thomisus spinifer</i> reared on immatures of <i>Rhopalosiphum maidis</i> at 27± 2°C.	138
28	Total protein, carbohydrates and lipids of adult and immature <i>Aphis craccivora</i> .	140
29	Total protein, carbohydrates and lipids of adult and immature <i>Rhopalosiphum maidis</i> .	142

LIST OF FIGURES

No.	Figure title	Page
1a	Mean temperature and percent relative humidity at Qaha station, Qalubya governorate during the broad bean season 2017.	54
1b	Seasonal dynamics of <i>A. craccivora</i> life stages infesting the broad bean crop during 2017 at Qaha station, Qalubya governorate.	54
1c	Seasonal dynamics of <i>Aphis craccivora</i> and prevalence of life stages collected on broad bean crop at Qaha station in 2017.	55
2a	Mean temperature and percent relative humidity at Qaha station, Qalubya governorate during the broad bean season 2018.	61
2b	Seasonal dynamics of <i>A. craccivora</i> life stages infesting the broad bean crop during 2018 at Qaha station, Qalubya governorate.	61
2c	Seasonal dynamics of <i>Aphis craccivora</i> and prevalence of life stages collected on broad bean crop at Qaha station in 2018.	62
3a	Mean temperature and percent relative humidity at Qaha station, Qalubya governorate during the maize season 2017.	68
3b	Seasonal dynamics of <i>R. maidis</i> life stages infesting the maize crop during 2017 at Qaha station, Qalubya governorate.	68

No.	Figure title	Page
3c	Seasonal dynamics of <i>Rhopalosiphum maidis</i> and prevalence of life stages collected on maize crop at Qaha station in 2017.	69
4a	Mean temperature and percent relative humidity at Qaha station, Qalubya governorate during the maize season 2018.	75
4b	Seasonal dynamics of <i>R. maidis</i> life stages infesting the maize crop during 2018 at Qaha station, Qalubya governorate.	75
4c	Seasonal dynamics of <i>Rhopalosiphum maidis</i> and prevalence of life stages collected on maize crop at Qaha station in 2018.	76
5	Number of collected spiders and percent occurrence of spider families on broad bean crop at Qaha station, Qalubya governorate during 2017 season	81
6	Number of collected spiders and percent occurrence of spider families on broad bean crop at Qaha station, Qalubya governorate during 2018 season.	81
7	Seasonal dynamics of spider families on broad bean crop at Qaha station, Qalubya governorate during 2017 season.	83
8	Seasonal dynamics of spider families on broad bean crop at Qaha station, Qalubya governorate during 2018 season.	85

No.	Figure title	Page
9	Number of collected spiders and percent occurrence of spider families on maize crop at Qaha station, Qalubya governorate during 2017 season.	89
10	Number of collected spiders and percent occurrence of spider families on maize crop at Qaha station, Qalubya governorate during 2018 season.	89
11	Seasonal dynamics of spider families on maize crop at Qaha station, Qalubya governorate during 2017 season.	91
12	Seasonal dynamics of spider families on maize crop at Qaha station, Qalubya governorate during 2018 season.	93
13a	Mean temperature and percent relative humidity at Qaha station, Qalubya governorate during the broad bean season 2017.	97
13b	Population density of spiders and their prey of <i>Aphis craccivora</i> on broad bean crop at Qaha station, Qalubya governorate during 2017 season.	97
14a	Mean temperature and percent relative humidity at Qaha station, Qalubya governorate during the broad bean season 2018.	98
14b	Population density of spiders and their prey of <i>Aphis craccivora</i> on broad bean crop at Qaha station, Qalubya governorate during 2018 season.	98
15a	Mean temperature and percent relative humidity at Qaha station, Qalubya governorate	100

No.	Figure title	Page
	during the maize season 2017.	
15b	Population density of spiders and their prey of <i>R. maidis</i> on maize crop at Qaha station, Qalubya governorate during 2017 season.	100
16a	Mean temperature and percent relative humidity at Qaha station, Qalubya governorate during the maize season 2018.	101
16b	Population density of spiders and their prey of <i>R. maidis</i> on maize crop at Qaha station, Qalubya governorate during 2018 season.	101
17a	Duration of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Aphis craccivora</i> at $27\pm 2^{\circ}\text{C}$.	114
17b	Food consumption of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Aphis craccivora</i> at $27\pm 2^{\circ}\text{C}$.	115
17c	Predation rate of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Aphis craccivora</i> at $27\pm 2^{\circ}\text{C}$.	116
18a	Duration of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Aphis craccivora</i> at $35\pm 2^{\circ}\text{C}$.	118
18b	Food consumption of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Aphis craccivora</i> at $35\pm 2^{\circ}\text{C}$.	119

No.	Figure title	Page
18c	Predation rate of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Aphis craccivora</i> at 35± 2°C .	120
19a	Duration of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 27± 2°C	122
19b	Food consumption of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 27± 2°C .	123
19c	Predation rate of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 27± 2°C .	124
20a	Duration of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 35± 2°C .	126
20b	Food consumption of the spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 35± 2°C.	127
20c	Predation rate of spiderlings of <i>Thanatus albini</i> (F: Philodromidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 35± 2°C .	128
21a	Duration of the spiderlings of <i>Thomisus spinifer</i> (F: Thomisidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 27± 2°C.	132

No.	Figure title	Page
21b	Food consumption of the spiderlings of <i>Thomisus spinifer</i> (F: Thomisidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 27± 2°C.	133
21c	Predation rate of the spiderlings of <i>Thomisus spinifer</i> (F: Thomisidae) fed on immatures and adult females of <i>Rhopalosiphum maidis</i> at 27± 2°C .	134
22	Total protein, carbohydrates and lipids of adult and immature <i>Aphis craccivora</i> .	141
23	Total protein, carbohydrates and lipids of adult and immature <i>Rhopalosiphum maidis</i> .	143