

MENOUFIA UNIVERSITY
Faculty of Agriculture
Department of Soil Science



RESPONSE OF SOME LEGUMINOUS PLANTS TO SPECIFIC RHIZOBIAL INOCULATION TIMING WITHIN GERMINATION STAGE

A Thesis Submitted

By

Bakre Abd Elgalel Mohamed Abd El-WAHAB

B.Sc. Agric. Sci., "Soils", Fac. of Agric., Suez Canal Univ. 1999
MSc. Soils, Fac. of Agric., Menoufia Univ. (2014)

As

A Partial Fulfillment of the Requirements

For

The Degree of
DOCTOR OF PHILOSOPHY

In

Agricultural Sciences "Soils"

Department of Soil Science

Menoufia University

2021

CONTENTS

SUBJECT	Page No.
Front Pages	(I-IV)
Front Sheet.....	i
Credit Sheet (Supervision Committee)	ii
Approval Sheet (arbitration Committee).....	iii
Acknowledgments.....	iv
1.INTRODUCTION.....	(1-3)
2. REVIEW OF LITERATURES	(4-21)
2.1. Biological Nitrogen Fixation	4
2.2. Rhizobia - Legume Symbiosis	6
2.3. Formulations	9
2.3.1. Liquid formulations	9
2.3.2. Solid formulations	10
2.4. Mechanisms of Rhizobia Plant Growth Promotion.	13
2.4.1. Direct mechanisms of Growth Promotion	14
2.4.2. Indirect mechanisms of growth promotion	14
2.5. Factors affecting biomass and nodulation	16
2.6. Rhizobial interaction for enhancing crop production	16
2.7. Rhizobium -Based Commercially Available Inoculants	19
2.8. Future Perspectives	20
3. MATERIAL AND METHODS	(22-38)
3.1. layout	22
3.1.1. Rhizobial specific strains	23
3.2. Some characteristics of the specific rhizobial strains	23
3.2.1. Morpho-cultural characterization	23
3.2.2. Temperature tolerance	25
3.2.3. Salt (NaCl) tolerance	25
3.2.4. Intrinsic sodium azide resistant	25
3.2.5. Resistance to lead (Pb)	25
3.2.6. Intrinsic antibiotic resistance	26
3.2.7. Indole acetic acid (IAA) production	26
3.2.8. Exopolysaccharide "EPS" production	27
3.2.9. Transmission electron microscopy	27

3.3. Application Experiment	28
3.3.1. Soil location and soil sampling	28
3.3.2. Plants	28
3.3.3. Inoculum preparations of the rhizobial specific strains	28
3.3.4. Liquid inoculation process	30
3.3.5. Solid inoculation process	30
3.3.6. Acetylene Reduction by Detached Acetylene Reduction by Detached Nodules ($\mu\text{mol C}_2\text{H}_4 / \text{gm dry nodule /hr.}$)	30
3.4. Methods	31
3.4.1. The experimental treatments	31
3.4.2. Field Experiment	33
3.5. Laboratory Determinations	35
3.5.1. Soil analysis	35
3.5.2. Plant analysis	37
3.6. Statistical Analysis	38
3.7. Media	38
4. RESULTS AND DISCUSSION (39- 136)	
4.1. Characterization of The Specific Rhizobial Strains	39
4.1.1. Morpho-Cultural Characteristics	39
4.1.2. Survival of the rhizobial specific strains at different temperature	41
4.1.3. Survival of the rhizobial specific strains at different salinity (NaCl) levels	41
4.1.4. Intrinsic sodium azide resistant	44
4.1.5. Resistance to lead (Pb)	46
4.1.6. Intrinsic antibiotic resistance	46
4.1.7. Indole acetic acid (IAA) production	50
4.1.8. Exopolysaccharide "EPS" (mg L^{-1}) production	53
4.1.9. Transmission electron microscopy	56
4.1.10. Interpreted characterization of the tested specific Rhizobial and Bradyrhizobial Strains	58
4.2. Responses of The Nodulation Process, Plant Growth and Crop Quality of Soybean, Common Bean and Peanut Plants to The Timing Application of Their Specific Rhizobial Liquid Inoculation	60
4.2.1. Effect of inoculation treatments on soybean plants	60
4.2.1.1. Number and weight of nodules	60
4.2.1.2. Nitrogenase activity	64
4.2.1.3. Fresh (FMY) and dry matter yields (DMY)	67
4.2.1.4. Macronutrient (N, P, K) contents	71
a- Nitrogen (N) contents	71
b-Phosphorus (P) contents	74
c-Potassium (K) contents	74

4.2.1.5. Micronutrient (Fe, Mn and Zn) contents	75
a- Iron (Fe) content	75
b- Zinc (Zn) content	77
c- Manganese (Mn) content	77
4.2.1.6. Grain and straw yields	78
4.2.1.7. Grain content of macronutrients	80
a- Nitrogen contents	80
b-Phosphorus (P) contents	82
c-Potassium (K) contents	83
4.2.1.8. Grain content of micronutrients	84
a- Iron (Fe) content	84
b- Zinc (Zn) content	86
c- Manganese (Mn) content	86
4.2.2. Effect of inoculation treatments on common bean plants	88
4.2.2.1. Number and weight of nodules	88
4.2.2.2. Nitrogenase activity	91
4.2.2.3. Fresh (FMY) and dry matter yields (DMY) of shoots common bean plants	93
4.2.2.4. Macronutrients (N, P, K) contents	96
a- Nitrogen (N) contents	96
b-Phosphorus (P) contents	99
c-Potassium (K) contents	99
4.2.2.5. Micronutrient (Fe, Mn, Zn) contents	100
a- Iron (Fe) content	100
b- Zinc (Zn) content	102
c- Manganese (Mn) content	102
4.2.2.6. Grain and straw yields	103
4.2.2.7. Grain content of macronutrients	105
a- Nitrogen contents	105
b-Phosphorus (P) contents	107
c-Potassium (K) contents	108
4.2.2.8. Grain content of micronutrients	108
a- Iron (Fe) content	108
b- Zinc (Zn) content	110
c- Manganese (Mn) content	111
4.2.3. Effect of inoculation treatments on peanut plants	112
4.2.3.1. Number and dry weight of nodules	112
4.2.3.2. Nitrogenase activity	115
4.2.3.3. Fresh (FMY) and dry matter yields (DMY) of shoots soybean plants	117
4.2.3.4. Macronutrient (N, P, K) contents	120
a- Nitrogen(N) contents	120
b-Phosphorus (P) contents	123
c-Potassium (K) contents	124
4.2.3.5. Micronutrient (Fe, Mn, Zn) contents	124

a- Iron (Fe) content	124
b- Zinc (Zn) content	126
c- Manganese (Mn) content	126
4.2.3.6. Grain and straw yields	127
4.2.3.7. Grain content of macronutrients	129
a- Nitrogen contents	129
b-Phosphorus (P) contents	131
c-Potassium (K) contents	131
4.2.3.8. Grain content of micronutrients	132
a- Iron (Fe) content	133
b- Zinc (Zn) content	135
c- Manganese (Mn) content	136
4.3. General Discussion	137- 142
5. SUMMARY	
	143- 157
6. REFERENCES	
	158- 172
ARABIC SUMMARY	
	1-10

List of Tables

Tables No.	Tables	Page No.
(1)	Physical and chemical properties and nutrient contents of the soil in the field experiment at Om Saber-Tahrir regions-Behera Governorate, Egypt.	29
(2)	Morpho-Cultural Characteristics of the tested specific rhizobial strains	40
(3)	Survival of the rhizobial specific strains at different temperature	42
(4)	Survival of the rhizobial specific strains at different salinity (NaCl) levels	43
(5)	Intrinsic tolerances of the rhizobial specific strains at different levels of sodium azide	45
(6)	Survival of the tested rhizobial specific strains at different Lead (Pb) concentrations.	47
(7)	Inhibition zone (mm) of the selected rhizobial strains affecting by some antibiotics additions	48
(8)	Indole acetic acid (IAA) ($\mu\text{g ml}^{-1}$) production by Rhizobial and Bradyrhizobial Strains	51
(9)	Exopolysaccharide "EPS" (mg L^{-1}) production by the specific Rhizobial and Bradyrhizobial Strains	54
(10)	The selected three strains that, represent the most efficient of the tested specific Rhizobial and Bradyrhizobial Strains	59
(11)	Nodule number (Nodule plant ⁻¹) and its dry weight (mg plant ⁻¹) and their RC values (%) of Giza 111 and Giza 135 cultivars soybean plants, at 45 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	62
(12)	Nitrogenase activity ($\mu\text{mol C}_2\text{H}_4/\text{gm dry nodule/hr}$) in nodule of Giza 111 and Giza 135 cultivars soybean root plants and their RC values %, at 45 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	66
(13)	Fresh and dry matter yields (FMY) (DMY) (Mg fed ⁻¹) and their RC values % of Giza 111 and Giza 135 cultivars soybean plants, at 60 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	68
(14)	Nitrogen (N), phosphorus (P) and potassium (K) concentration (%) and uptake (Kg fed ⁻¹) in shoots (60 days after sowing) and their RC values % for Giza 111 and Giza 135 cultivars soybean plants, as affected by the timing application of its specific rhizobial liquid inoculation.	74
(15)	Iron (Fe), Manganese (Mn) and Zinc (Zn) concentration (mg Kg ⁻¹) and uptake (g fed ⁻¹) in shoots (60 days after sowing) and their RC values % for Giza 111 and Giza 135 cultivars soybean plants, as affected by the timing application of its specific rhizobial liquid inoculation.	76
(16)	Grain and straw yields (Kg fed ⁻¹) and their RC values % of Giza 111 and Giza 135 cultivars soybean plants, at harvest, as affected by the timing application of its specific rhizobial liquid inoculation.	79

(17)	Nitrogen (N), phosphorus (P) and potassium (K) concentration (g Kg ⁻¹) and uptake (Kg fed-1) and their RC values % in grain of Giza 111 and Giza 135 cultivars soybean plants, as affected by the timing application of its specific rhizobial liquid inoculation.	81
(18)	Iron (Fe), Manganese (Mn) and Zinc (Zn) concentration (mg Kg ⁻¹) and uptake (g fed-1) in grains and their RC values % of Giza 111 and Giza 135 cultivars soybean plants, as affected by the timing application of its specific rhizobial liquid inoculation.	86
(19)	Nodule number (Nodule plant ⁻¹) and its dry weight (mg plant ⁻¹) and their RC values % of Giza 4 and Giza 6 cultivars common bean plants, at 45 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	89
(20)	Nitrogenase activity (μmol C ₂ H ₄ /gm dry nodule/hr.) in nodule on Giza 4 and Giza 6 cultivars common bean root plants and their RC values %, at 45 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	92
(21)	Fresh and dry matter yields (FMY) (DMY) (Mg fed-1) and their RC values % of Giza 4 and Giza 6 cultivars common bean plants, at 60 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	94
(22)	Nitrogen (N), phosphorus (P) and potassium (K) concentration (%) and uptake (Kg fed-1) in shoots (60 days after sowing) and their RC values % for Giza 4 and Giza 6 cultivars common bean plants, as affected by the timing application of its specific rhizobial liquid inoculation.	97
(23)	Iron (Fe), Zinc (Zn) and Manganese (Mn) concentration (mg Kg ⁻¹) and uptake (g fed-1) in shoots (60 days after sowing) and their RC values % for Giza 4 and Giza 6 cultivars common ben plants, as affected by the timing application of its specific rhizobial liquid inoculation.	101
(24)	Grain and straw yields (Kg/fed) and their RC values % of Giza 4 and Giza 6 cultivars common bean plants, at harvest, as affected by the timing application of its specific rhizobial liquid inoculation.	104
(25)	Nitrogen (N), phosphorus (P) and potassium (K) concentration (g Kg ⁻¹) and uptake (Kg fed-1) and their RC values % in grains of Giza 4 and Giza 6 cultivars common bean plants, as affected by the timing application of its specific rhizobial liquid inoculation.	106
(26)	Iron (Fe), Manganese (Mn) and Zinc (Zn) concentration (mg Kg ⁻¹) and uptake (g fed-1) in grains and their RC values % of Giza 4 and Giza 6 cultivars common bean plants, as affected by the timing application of its specific rhizobial liquid inoculation.	109
(27)	Nodule number (Nodule plant ⁻¹) and its dry weight (mg plant ⁻¹) and their RC values % of Ponch and NC cultivars peanut plants, at 45 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	113
(28)	Nitrogenase activity (μmol C ₂ H ₄ / gm dry nodule/hr.) in nodule on Ponch and NC cultivars peanut root plants and their RC values %, at 45 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	116
(29)	Fresh and dry matter yields (FMY) (DMY) (Mg fed-1) and their RC	118

	values % of Ponch and NC cultivars peanut plants, at 60 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	
(30)	Nitrogen (N), phosphorus (P) and potassium (K) concentration (%) and uptake (Kg fed-1) in shoots (60 days after sowing) and their RC values % for Ponch and NC cultivars peanut plants, as affected by the timing application of its specific rhizobial liquid inoculation.	122
(31)	Iron (Fe), Manganese (Mn) and Zinc (Zn) concentration (mg Kg-1) and uptake (g fed-1) in shoots (60 days after sowing) and their RC values % for Ponch and NC cultivars peanut plants, as affected by the timing application of its specific rhizobial liquid inoculation.	125
(32)	Grain and straw yields (Kg fed-1) and their RC values % of Ponch and NC cultivars peanut plants, at harvest, as affected by the timing application of its specific rhizobial liquid inoculation.	128
(33)	Nitrogen (N), phosphorus (P) and potassium (K) concentration (g Kg-1) and uptake (Kg fed-1) and their RC values % in grains of Ponch and NS cultivars peanut plants, as affected by the timing application of its specific rhizobial liquid inoculation.	130
(34)	Iron (Fe), Zinc (Zn) and Manganese (Mn) concentration (mg Kg-1) and uptake (g fed-1) in grains and their RC values % of Ponch and NC cultivars peanut plants, as affected by the timing application of its specific rhizobial liquid inoculation.	134

List of Diagram

Figures NO.	Figures	Page NO.
(1)	Source of Bradyrhizobium and Rhizobium strains and isolates used in the lab and field experiments	24

List of Figures

Figures NO.	Figures	Page NO.
(1)	The experimental design (for one replicate) and Applying treatments.	32
(2)	Indol acitic acid (IAA) ($\mu\text{g ml}^{-1}$) production by Rhizobial and Bradyrhizobial Strains	52
(3)	Exopolysaccharide "EPS" (mg L^{-1}) production by the specific Rhizobial and Bradyrhizobial Strains	55
(4)	Transmission electron micrographs of: (a) Rhizobium leguminosarum strain (203) showing monotrichous flagella (b) Bradyrhizobium SP strain (617) showing peritrichous flagellium. (c) Rhizobium phaseoli strain (301) showing amphitrichous flagellium. (d) Bradyrhizobium japonicum strain (3307) showing lophotrichous flagelium. Scale bar: $0.5\mu\text{m}$.	57
(5)	Fresh and dry matter yields (FMY) (DMY) (Mg fed^{-1}) and their RC values % of Giza 111 and Giza 135 soybean plants cultivar, at 60 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	69
(6)	Fresh and dry matter yields (FMY) (DMY) (Mg fed^{-1}) and their RC values % of Giza 4 and Giza 6 cultivars common bean plants, at 60 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	95
(7)	Fresh and dry matter yields (FMY) (DMY) (Mg fed^{-1}) and their RC values % of Ponch and NC cultivars peanut plants, at 60 days after sowing, as affected by the timing application of its specific rhizobial liquid inoculation.	119

5. SUMMARY

The present work was conducted to: a) Evaluation the effect of some abiotic stress tolerance of the specific rhizobial strains and its phenotypic characterization. b) Evaluation the effect of the timing application of the specific rhizobial liquid inoculation to some leguminous plants on the nodulation process. c) Evaluation the response of some hybrids of the same leguminous plants to the timing application of its specific rhizobial liquid inoculation., and d) Reflection between the response of some leguminous plants and their hybrids to the timing application of its specific rhizobial liquid inoculation on the plant growth and crop quality.

To select the most three efficient strains to employed in the field experiment. Some specific rhizobial strains were subjected to some abiotic stress tolerance and its phenotypic characterization. Antibiotics, temperature, salt (NaCl) lead (Pb) and sodium azide tolerance, IAA, exopolysaccharides (EPS), were determined, as well as morpho-cultural characterization (Gram staining, colony size, shape, border, elevation, color, mucosity, transparency, scanning electron micro graph of the same tested specific rhizobial strains, were performed.

Field Experiment

Three field experiments were conducted at the same site at Om Saber Village-Tahrir regions - Beheira Governorate, Egypt (Latitude 30°40'N- Longitude 30°33'E) (sandy soils presented newly reclaimed soil). The first, second and third experiments were carried out on soybean "(*Glycine max*), common bean plants (*Phaseolus vulgaris L.*) and peanut (*Arachis hypogaea L.*), respectively. Each experiment was carried using two cultivars of the cultivated leguminous plants. These treatment of each experiments were

complete randomized block plot design. The planting dates were at 15 April and 15 Mai, 1st September 2019 and for Soybean, common and peanut bean plants, respectively.

Rhizobial specific strains

In this study, the used three rhizobial specific strains were selected among other specific strains, which were subjected to some abiotic stress tolerance and its phenotypic characterization. These strains were: *Rhizobium legumseram var phaseoli*, *Bradyrhizobium japonicum* and *Bradyrhizobium spp* strains for inoculation seeds of common bean (*Phaseolus vulgaris*L.), Soybean (*Glycine max*) and peanut (*Arachis hypogaea* L.), respectively. These strains were kindly provided from Lab of Microbiology, Soil, Water and Environment Research Institute (SWERI), Agriculture Research Center (ARC), Giza, Egypt.

Responses of the nodulation process, plant growth and crop quality to the timing application of their specific rhizobial liquid inoculation was examined in a separately three field experiments. For this propose, the three leguminous plant seeds were continuously inoculated with their specific rhizobial at 0-7th (First Week "FW"), 8-14th (Second Week "SW") and 15-21th (third week "TW") days after sowing, through drip irrigation system. Indeed, in these experiments the solid inoculum was applied (at the sowing) as a reference for the liquid inoculation.

a. Inoculum preparations of the rhizobial specific strains.

Inoculum for these experiments was obtained by growing the specific rhizobial strains in **Yeast Mannitol Broth (YMB)** medium. All strains were maintained in **YMB** and cultured routinely at 28°C.

b. Liquid inoculation process

All plots of liquid inoculation of each sub treatment were continuously inoculated with its specific rhizobial strain at 0-7th (First Week "FW"), 8-14th (Second Week "SW") and 15-21th (third week "TW") days after sowing, through the dripping system.

c. Solid inoculation (SI) process

All plots of solid inoculation of each sub treatment were inoculated with its solid specific rhizobial strains (Okadin) just before sowing. The solid inoculation were performed by coating the seeds at a rate of 800 gm of solid inoculum/fed.. Three different bags of Biofertilizer (Okadin) (each bag was specific for each leguminous crop).

The experimental treatments

The experimental treatments in this study were arranged within the experimental units in split randomized block design in three replicates illustrated in table (1) as follows:

a-Main treatments:

Five main treatments: Control treatment (without any inoculation), solid inoculation "SI" at sowing and liquid inoculation which carried out through (via drip irrigation "Fertigation") at three times i.e. at 0-7th (First Week "FW"), 8-14th (Second Week "SW") and 15-21th (Third Week "TW") days after sowing, through the dripping system, were used in this study.

b- Sub-treatments:

These sub treatments represented by the cultured three leguminous plants These plants were sub treatments soybean plants "(Glycine max) "Giza 111 and Giza 135 cultivars" common bean plants (*Phaseolus vulgaris L.*) "Giza 4 and Giza 6 cultivars", and peanut plants (*Arachis hypogaea L.*) "Ponch and NC" Cultivars.

c- Sub sub treatment:

These sub sub treatments were represented by two cultivars of the cultivated crops.

The obtained results could be summarized as follows:

I. Selected three strains that, represent the most efficient of the tested specific Rhizobial and Bradyrhizobial Strains

Some specific rhizobial strains were subjected to some abiotic stress tolerance and its phenotypic characterization, to select the most three efficient strains to employed in the field experiment. These three specific rhizobial strains showed, higher potentials of IAA and EPS production, high survival capabilities at different temperature degrees, salinity levels, lead (Pb), intrinsic sodium azide resistant, intrinsic antibiotic resistance in compared to the other tested specific Rhizobial and Bradyrhizobial strains.

II. . Responses of The Nodulation Process, Plant Growth and Crop Quality of Soybean, Common Bean and Peanut Plants to The Timing Application of Their Specific Rhizobial Liquid Inoculation

A . Soybean Plants

1. Effect of inoculation treatments on nodulation

- The maximum number, weight nodules and nitorgenase activity were observed with the *B. japonicum* liquid application through the first and at the second week "FW" and "SW" after sowing compared to the other liquid or solid inoculation treatments.
- The correspondence values of number and weight nodules were 50, 68, 65 and 48 nodule plant⁻¹ and 13.05, 42.30, 32.60 and 12,42 mg plant⁻¹ for the "SI", "FW", "SW", "SW", and "TW", after sowing, for the Giza 111 cultivar. whereas, for the Giza 135 cultivar were 64.00, 76.00, 75.00

and nodule plant⁻¹ and 13.67, 51.33, 33.08 and 13.12 mg plant⁻¹, with the same abovementioned treatments.

- All values of RC of weight nodules and nitrogenase activity were higher under the treatments of the *B. japonicum* liquid applied at the first week "FW" after sowing, with both the two soybean cultivars, than those found with other studied experimental treatments.

2. Fresh and dry matter yield

- The obtained fresh and dry matter yield of Giza 135 soybean cultivar plants were higher than those found of Giza 111 cultivar with all studied inoculation treatments.
- The value of shoots FMY and DMY achieved; 6.483 and 3.848 Mg fed⁻¹ for the plants of Giza 135 inoculated with the liquid *B. japonicum* at "FW" after sowing, respectively, in compared to 4.078 and 2.305 Mg fed⁻¹ for FMY and DMY of Giza 111 soybean cultivar plants, under the same abovementioned liquid inoculation treatment, respectively.

3. Macronutrients contents in shoots of soybean plants.

- It was noticed that nitrogen content (%) and its uptake as Kg fed⁻¹ in shoots were significantly enhanced by the liquid inoculation by *B. japonicum* at the first "FW" and at the second "SW" week in compared to the solid inoculation (SI) and the liquid inoculation by *B. japonicum* at the third week (TW).
- In regarding to Giza 111 and Giza135 cultivar soybean plants, inoculated with liquid *B. japonicum* 3307 at the first week "FW" after sowing, led to maximum values of P uptake by shoots which was: 15.37 and 26.55 Kg fed⁻¹, respectively.
- The inoculation treatments with liquid *B. japonicum* 3307 at the first "FW" and at the second "SW" week after sowing, significantly

augmented K uptake in shoots of soybean plants compared to the other solid "SI" or liquid inoculation treatments "TW".

4. Micronutrient contents in shoots of soybean plants.

- The highest values of Fe content in shoots were: 1080.67 and 707.83 and 1147.67 and 747.00 mg Kg⁻¹ which equal 2490.94 and 1424.25 and 4416.25 and 2699.13 g fed⁻¹, were found in shoot plants inoculated by liquid inoculation at "FW" and "SW" after sowing, of Giza 111 and Giza 135, respectively.
- The highest promotion of Zn and Mn contents were found in the soybean shoot plants received the liquid inoculum by *B. japonicum* at the first "FW" and the second "SW" week after sowing in Giza 111 and Giza 135 cultivar plants.

5. Grain and straw yield of soybean plants.

- Inoculation with liquid *B. japonicum* treatments at "FW" and "SW" week after sowing, gained a higher grain yields: 2784.6 and 2687.64 Kg fed⁻¹ of Giza 111. Whereas, the grain and straw yields of Giza 135 cultivars soybean plants were: 2939.3 and 2867.9 Kg fed⁻¹ respectively, with the same aforesaid liquid inoculation treatments.

6. Grain content of macronutrients

- Liquid applications of *B. japonicum* at "FW" and "SW" after sowing resulted in a maximum increase in the grain content of N with the Giza 135 it were: 62.77 and 62.77 g Kg⁻¹ and 184.64 180.13 Kg fed⁻¹, compared to the 58.77 57.77 g Kg⁻¹ and 163.8 and 160.98 Kg fed⁻¹ with the Giza 1112.
- The highest grains P and K contents were found in the plants inoculated with the liquid *B. japonicum* at "FW" and "SW", after sowing.

compared to the solid inoculation "SI" and the liquid *B. japonicum* at "TW" treatments.

7. Grain content of micronutrients

- The plants successive received the liquid *B. japonicum* for all first week "FW" and second "SW" week after sowing, attained higher values of grain Fe concentration and its uptake by 712.98 and 711.98 mg Kg⁻¹ which equal 1985.37 and 1982.58 g fed⁻¹ of Giza 111, respectively. Whereas, it were 764.98 and 721.98 mg Kg⁻¹ which equal 2248.5 and 2070.57 g fed⁻¹ of Giza 135, respectively.

B. Common Bean Plants

1. Effect of inoculation treatments on nodulation

- The maximum number and weight nodules were observed with the *R. phaseoli* liquid application the first "FW" and the second "SW" week after sowing compared to the later application of liquid *R. phaseoli* or the solid inoculation treatments.
- The correspondence values of number and weight nodules were 68.00 and 65.00 nodule plant⁻¹ which equal 517.83 and 502.17 mg plant⁻¹ for the Giza 4 cultivar. whereas, for the Giza 6 cultivar were 75.00 and 74.00 nodule plant⁻¹ which equal 776.67 and 688.67mg/plant, with the same treatments of *R. phaseoli* liquid application.
- The data showed in generally increases in nitorgenase activity (symbiotic effectiveness) of nodules formed on common bean root plants inoculated with liquid *R. phaseoli* at the first "FW" and the second "SW" week after sowing, in both common bean cultivar plants compared to the other applied inoculation treatments.

2. Fresh and dry matter yield

- These data show that, the treatments of inoculation with the *R. phaseoli liquid* application at "FW" after sowing was associated by an increase of fresh (FMY) and dry (DMY) shoots of both studied common bean cultivar plants.
- The obtained fresh and dry matter yield of Giza 6 common bean cultivar plants were higher than those found of Giza 4 cultivar with all studied inoculation treatments.
- The higher nodulation and accumulation of shoot biomass promoted by the liquid inoculation by *R. phaseoli* at first "FW" and "FW" week after sowing, indicate that these timing applications are promising inoculation methods at the early germination stages of development for common bean growth and productivity.

3. Macronutrient (N, P, K) contents of common bean plants.

- The inoculation treatment with the liquid *R. phaseoli* a at "FW" and "SW" week after sowing, induced the maximum values of nitrogen content (%) and Kg fed⁻¹ of both cultivars (Giza 4 and Giza 6).
- Nitrogen content (%) and its uptake as Kg fed⁻¹ in shoots were significantly enhanced by the early application of liquid inoculation by *R. phaseoli* ("FW" and "SW") in compared to the late application of the same liquid *Rhizobia* at "TW" or solid inoculation "SI".
- The inoculated plants with liquid *R. phaseoli* 301 for "FW" and "SW" week after sowing appeared a high P concentration, in shoots of the two cultivar common bean plants. In regarding to Giza 4 and Giza 6 cultivar common bean plants, inoculated with liquid *R. phaseoli* 301 at "FW" and "SW" week after sowing, led to

maximum values of P uptake by shoots which was: 6.03 and 5.63 and 9.75 Kg fed⁻¹ and 9.49 Kg fed⁻¹, respectively.

4. Micronutrient contents in shoots of common bean plants.

- The liquid inoculation with *R. phaseoli* at "**FW**" and "**SW**", led to a significant increases of Fe content and its uptake in shoot of common bean plants compared to the "**SI**" and the liquid inoculation with *R. phaseoli* at "**TW**".
- Moreover, the highest values of Fe content in shoots were: 1950.50 and 2107.67 mg kg⁻¹ which equal 2901.75 and 3224.50 Kg fed⁻¹, were observed with treatments of liquid inoculation at "**FW**" of Giza 4 and Giza 6, respectively.
- The highest promotion of Zn and Mn content were found in the common bean shoot plants received the liquid inoculum by *R. phaseoli* for successive 7 days during the first ("**FW**") week after sowing, in Giza 4 and Giza 6 cultivar plants.

5. Grain and straw yield of common bean plants.

- The inoculation treatments by solid or liquid *R. phaseoli* 301 significantly augmented grain and straw yields of Giza 4 and Giza 6 cultivars common bean plants, compared to the uninoculated one. Inoculation with liquid *R. phaseoli* treatments at the "**FW**" and "**FW**" week after sowing, gained a significantly higher grain yields: 1401.40 and 1373.4 Kg fed⁻¹ and 1684.2 and 1661.1 Kg fed⁻¹ of Giza 4 and Giza 6 cultivar common bean plants, respectively.
- Similar increase effect of liquid inoculation "**FW**" and "**FW**" treatments was found with straw yields of the examined common bean cultivar plants. It is clear from of Giza 6 cultivar showed the

highest significantly values of grain and straw yields, compared those with Giza 4 cultivar, under all treatments studied.

- In respect to RC %, values of inoculation treatments by liquid *R. phaseoli* at the "FW" and "FW", gave the highest RC (%) values of grain yields: 67,25 and 63.91% of Giza 4 cultivar, and 62.02 and 59.80% of Giza 6 cultivar, respectively.

6. Grain content of macronutrients

- Data appeared that, there are widely varied according to the planted cultivars and type and applying times of the inoculum. For example, with all inoculation studied treatments N, P and K contents (g/Kg and Kg fed⁻¹) in the grains of Giza 6 cultivars was higher than that found in the grains of Giza 4 plants.
- Generally, applications of liquid inoculation by *R. phaseoli* at the "FW" and "FW" resulted in a significant increase the grains of common bean plants N, P and K contents (%) and its relative change "RC". These findings related with the effect of liquid inoculation with *R. phaseoli* at "FW" and "FW" weeks, after sowing, to the establishing the symbiotic relationship between the *R. phaseoli* and the lateral roots in the beginning of the germination stage of common bean plants.

7. Grain content of micronutrients

The data displayed that, the concentrations and uptake of the determined micronutrients, *i.e.* iron "Fe", zinc "Zn" and manganese "Mn" and the rates of their relative changes of their contents "RC, %", in the grains of two common bean cultivars plants grown on the new reclaimed sandy soils, as affected by inoculation with the solid or liquid *R. phaseoli* 3307.

C. Peanut Plants

1. Effect of inoculation treatments on nodulation

- The maximum number and weight nodules were observed with the liquid *Bradyrhizobium spp. 3339* application successively at the first week "FW" after sowing compared to the other liquid or solid inoculation "SI" treatments. The correspondence values of number and weight nodules were 37.33 nodule plant⁻¹ and 660.67 mg plant⁻¹ for. liquid *Bradyrhizobium spp* application at first "FW" after sowing, for the Ponch cultivar. whereas, for the NC cultivar were 40.0 nodule plant⁻¹ and 911.50 mg plant⁻¹, with the same treatments of *Bradyrhizobium spp. 3339* liquid application
- The RC (%) values of nitrogenase activity of nodules formed on peanut root plants received the liquid *Bradyrhizobium spp. 3339* in both peanut cultivar plants, attained higher values of RC of the nitrogenase activity compared with that received the "SI" and "TW" treatments.

2. Fresh and dry matter yield

- The value of shoots FMY and DMY achieved; 19.63 and 5.79 Mg fed⁻¹ for the plants of Ponch inoculated with the liquid *Bradyrhizobium spp.* at "FW", respectively, in compared to 28.61 and 11.37 Mg fed⁻¹ for FMY and DMY of NC peanut cultivar plants, under the same abovementioned liquid inoculation treatment, respectively.

3. Macronutrients contents in shoots of peanut plants.

- The inoculation treatment with the liquid *Bradyrhizobium spp. 3339* at "FW", and "SW" week after sowing, induced the maximum values of nitrogen content (%) and Kg fed⁻¹ of both cultivars (Ponch

and NC). Nitrogen contents of shoots were: 3.02 and 2.99%, which equal to 174.96 and 166.17 Kg fed⁻¹ of Ponch cultivar, increased to 3.54 and 3.23% which equal to 401.87 and 274.14 Kg fed⁻¹ of NC cultivar, respectively. It was noticed that nitrogen content (%) and its uptake as Kg fed⁻¹, in seeds were significantly enhanced by the liquid inoculation at "FW", and "SW" in compared to the "SI" or "TW" treatments.

- In regarding to Ponch and NC cultivar peanut plants, inoculated with liquid *Bradyrhizobium spp.* at "FW" week after sowing, led to maximum values of P uptake by shoots which was: 24.03 and 57.98 Kg fed⁻¹, respectively.
- The plants inoculated with the same inoculation treatments ("FW") gave a higher values of K contents in shoots *i.e.* 0.99 and 1.05 % which equal to 57.32 and 118.80 Kg fed⁻¹ of Ponch and NC cultivar plants, respectively.

4. Micronutrient contents in shoots of peanut plants.

- The highest values of Fe uptake by shoots were obtained with the liquid inoculation with *Bradyrhizobium spp.* at "FW" and "SW" week after sowing were: 3837.0 and 2077.3 and 9286.0 and 4202.8 g fed⁻¹ of Ponch and NC, respectively.
- The highest promotion of Zn and Mn content was found in the peanut shoot plants received the liquid inoculum by *Bradyrhizobium spp.* at "FW" week after sowing, in Ponch and NC cultivar plants.

5. Grain and straw yields (Kg fed⁻¹) of peanut plants.

- Inoculation with liquid *Bradyrhizobium spp.* 3339 treatments at "FW" and "SW" week after sowing, gained a higher grain yields: 1551.21 and 1438.49 Kg fed⁻¹ and 1682.17 and 1483.58 Kg fed⁻¹ of Ponch and NC cultivars peanut plants, respectively. Similar increase

effect of liquid inoculation treatments was found with straw yields of the examined peanut cultivar plants.

- It is clear that, NC cultivar showed the highest significantly values of grain and straw yields, compared those with Ponch cultivar, under all treatments studied.

6. Grain content of macro and micronutrients

- Data appeared that, there are widely varied according to the planted cultivars and type and applying times of the liquid inoculation.
- For example, with all inoculation studied treatments N, P and K content (g Kg^{-1} and Kg fed^{-1}) in the grains of NC cultivars was higher than that found in the grains of Ponch cultivar.
- Generally, applications of liquid inoculation by *Bradyrhizobium spp.* 3339 at "FW" and "SW" week after sowing, resulted in a significant increase the grains of peanut plants N content (%) and its relative change "RC".
- These findings related with the effect of liquid inoculation at "FW" and "SW" week after sowing, on the availability of soil N, P and K and its uptake by plants.
- Data recorded manifested that, the concentrations and uptake of the determined micronutrients, *i.e.* iron "Fe", zinc "Zn" and manganese "Mn" and the rates of their relative changes of their contents "RC, %", in the grains of two peanut cultivars plants grown on the new reclaimed sandy soils, were affected by inoculation with the solid or liquid *Bradyrhizobium spp.* 617.

General Conclusion

In these study, data pointed out that, the timing application of specific Rhizobial liquid inoculation (*Bradyrhizobium japonicum* 3307, *Rhizobium legumseram var phaseoli* 301 and *Bradyrhizobium spp.* 3339) of soyabean,

common bean and peanut plants, respectively, at the first "FW" and at the second "SW" week after sowing, were the most efficient inoculation treatments according to the high values of nodulation efficiency, macro and micronutrients (N, P and K& Fe Mn and Zn) concentrations and uptake by the three tested leguminous plants (high significant) and produced the highest fresh and dry matter yields at all experimental treatments. These findings are due to the successive application of liquid *B. japonicum* 3307, *R. phaseoli* 301 and *Bradyrhizobium* spp. 3339, at "FW" and at "SW" week after sowing, gave capability of leguminous plants form a symbiotic relationship (nodule formation) with a host-specific strains during their germination stage, consequently, further nodule development in newly developing root tissue. Meanwhile, the use of liquid inoculation at "FW" and at "SW" week after sowing, consequently adding huge number of rhizobia on the grains practically at the early germination stage were more effective competitive, and adapted strains to inoculate soybean, common bean and peanut plants, such as *the* specific rhizobial and *Bradyrhizobial* strains compared to the solid inoculation "SI" or the late liquid inoculation at "TW". It is important to mention that, when liquid *B. japonicum* 3307, *R. phaseoli* 301 and *Bradyrhizobium* spp. 3339, applications were applied at third "TW" week after sowing, The seeds had gone through the germination in earlier period and thus, did not affect the nodule numbers and did not affect the symbiotic effectiveness.

In additionally, in this study, drip irrigation allowed us to application of liquid of *B. japonicum* 3307, *R. phaseoli* 301 and *Bradyrhizobium* spp. 617, at the "FW", "SW" and "TW" week after sowing, consequently, these specific strains maybe play as plant growth promoting rhizobacteria (PGPR).

As an outlook at the results of the nodule number, nitrogenase activity, fresh and dry matter yield N, P, K, Fe, Mn and Zn contents in soyabean,

common bean and peanut shoot plants, and grains (**Tables 11 to 34**), the following orders are inferred, as affected by solid or liquid inoculation by of *B. japonicum* 3307, *R. phaseoli* 301 and *Bradyrhizobium spp.* 617:

- As for inoculation treatments:

Liquid Inoculation at "FW" > Liquid Inoculation at "SW" > solid Inoculation "SI" > Liquid Inoculation at "TW"

As for soybean cultivars:

Giza 135 > Giza 111 cultivar.

As for common bean cultivars:

Giza 6 > Giza 4 cultivar.

As for peanut cultivars:

NC > Ponch cultivar.

Further inoculation studies in leguminous plants with liquid inoculation within this hypothesis. the early 7 days immediately after sowing and responses of the nodulation process, plant growth and rop Quality during vegetative and reproductive developmental stages are needed to test .